

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
 United States Patent and Trademark
 Office
 Box PCT
 Washington, D.C.20231
 ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 03 October 2000 (03.10.00)	
International application No. PCT/JP00/01022	Applicant's or agent's file reference 661745
International filing date (day/month/year) 23 February 2000 (23.02.00)	Priority date (day/month/year) 03 March 1999 (03.03.99)
Applicant SATO, Fuminori et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

24 August 2000 (24.08.00)

☐ in a notice effecting later election filed with the International Bureau on:2. The election ☒ was☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland	Authorized officer Henrik Nyberg
Facsimile No.: (41-22) 740.14.35	Telephone No.: (41-22) 338.83.38

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

70485 lwo

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 307/68, C07C 255/29, C07D 333/24, C07K 5/062, 5/078, 5/065, C07C 323/60, 317/50, A61K 31/275, 31/277, 31/341, 31/381, A61P 11/00</p>	A1	<p>(11) International Publication Number: WO 00/49008</p> <p>(43) International Publication Date: 24 August 2000 (24.08.00)</p>								
<p>(21) International Application Number: PCT/GB00/00529</p> <p>(22) International Filing Date: 16 February 2000 (16.02.00)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>9903853.1</td> <td>20 February 1999 (20.02.99)</td> <td>GB</td> </tr> <tr> <td>9916099.6</td> <td>10 July 1999 (10.07.99)</td> <td>GB</td> </tr> <tr> <td>9917171.2</td> <td>23 July 1999 (23.07.99)</td> <td>GB</td> </tr> </table> <p>(71) Applicant (for all designated States except US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): TUCKER, Howard [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). LARGE, Michael, Stewart [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). OLDFIELD, John [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). JOHNSTONE, Craig [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB). EDWARDS, Philip, Neil [GB/GB]; Alderley Park, Macclesfield, Cheshire SK10 4TG (GB).</p>	9903853.1	20 February 1999 (20.02.99)	GB	9916099.6	10 July 1999 (10.07.99)	GB	9917171.2	23 July 1999 (23.07.99)	GB	<p>(74) Agent: TIERNEY, Francis, John; AstraZeneca, Global Intellectual Property, P.O.Box 272, Mereside, Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).</p> <p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p><i>With international search report.</i></p> <p><i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
9903853.1	20 February 1999 (20.02.99)	GB								
9916099.6	10 July 1999 (10.07.99)	GB								
9917171.2	23 July 1999 (23.07.99)	GB								
<p>(54) Title: DI- AND TRIPEPTIDE NITRILE DERIVATIVES AS INHIBITORS OF CATHEPSIN L AND CATHEPSIN S</p> <div style="text-align: center; margin: 20px 0;"> $\text{R}^1-(\text{AA}^1)_r-(\text{AA}^2)-\underset{\text{H}}{\text{N}}-\overset{\text{R}^2}{\text{C}}(\text{R}^3)-\text{CN} \quad (1)$ </div>										
<p>(57) Abstract</p> <p>A compound of formula (I), wherein R¹, R², R³, AA¹, AA² and r are defined; a composition comprising a compound of formula (I) and a carrier or diluent; a compound of formula (I) for use as a medicament; the use of a compound of formula (I) in the manufacture of a medicament for use in the inhibition of a cysteine protease in a warm blooded animal; the use of a compound of formula (I) in the manufacture of a medicament for use in the treatment of chronic obstructive pulmonary disease in a warm blooded animal; and a method of treating a Cathepsin L or Cathepsin S mediated disease state in mammals which comprises administering to a mammal in need of such treatment an effective amount of a compound of formula (I).</p>										

INTERNATIONAL SEARCH REPORT

Inter national Application No
PCT/GB 00/00529

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D307/68 C07C255/29 C07D333/24 C07K5/062 C07K5/078
C07K5/065 C07C323/60 C07C317/50 A61K31/275 A61K31/277
A61K31/341 A61K31/381 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D C07K A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. SUZUE, ET AL.: "Hepatic agents. I. Synthesis of aminoacyl (and hydroxyacyl) aminoacetonitriles" CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 16, no. 8, August 1968 (1968-08), pages 1417-1432, XP002108053 Pharmaceutical Society of Japan, Tokyo, JP ISSN: 0009-2363 the whole document --- -/--	1-4, 6, 7, 10-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *B* document member of the same patent family

Date of the actual completion of the international search

14 June 2000

Date of mailing of the international search report

30/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCT/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A.R. KATRITZKY, ET AL.: "Benzotriazole-assisted synthesis of alpha-aminonitriles and a conceptually novel method for peptide elongation" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 6, June 1990 (1990-06), pages 1853-1857, XP002140042 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X compounds 5g-5m, 6g-6m ---	1,2,4,7, 11
X	GB 1 073 288 A (KYORIN SEIYAKU) 21 June 1967 (1967-06-21) the whole document ---	1,7,10, 12-14
X	S.A. THOMPSON, ET AL.: "Carboxy-modified amino acids and peptides as protease inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 29, no. 1, January 1986 (1986-01), pages 104-111, XP002140043 American Chemical Society, Washington, DC, US ISSN: 0022-2623 compounds 3b, 3d, 4b, 4d ---	1,7, 10-12
X	D.T. ELMORE, ET AL.: "Thioesters of amino acid derivatives as thioacylating agents in thiopeptide synthesis" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 5, May 1988 (1988-05), pages 1051-1055, XP002140044 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X page 1051, right-hand column, line 10 - line 26 ---	1-3,6,7, 10
X	H. MOSER, ET AL.: "Poly(dipeptamidium)-Salze: Definition und Methoden zur präparativen Herstellung" HELVETICA CHIMICA ACTA, vol. 69, 1986, pages 1224-1262, XP002049723 Verlag Helvetica Chimica Aata, Basle, CH ISSN: 0018-019X compounds 3, 4a, 5 ---	1-3,6,7, 10-12
	-/--	

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCT/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	D.J. BUTTLE, ET AL.: "Affinity purification of the novel cysteine proteinase papaya proteinase IV, and papain from papaya latex" BIOCHEMICAL JOURNAL, vol. 261, no. 2, 15 July 1989 (1989-07-15), pages 469-476, XP000913659 Portland Press, London, GB ISSN: 0264-6021 page 471, right-hand column	1
X	S. LIU, ET AL.: "Effects of ligand homologation and ligand reactivity on the apparent kinetic specificity of papain" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1250, 1995, pages 43-48, XP002140045 Elsevier Science, Amsterdam, NL ISSN: 0006-3002 compound 3a	1
X	C.A. LEWIS, ET AL.: "Thiohemiacetal formation by inhibitory aldehydes at the active site" BIOCHEMISTRY, vol. 16, no. 22, 1 November 1977 (1977-11-01), pages 4890-4895, XP002140046 American Chemical Society, Washington, DC, US ISSN: 0006-2960 table II, entries 8, 11	1
P, X	WO 99 24460 A (NOVARTIS) 20 May 1999 (1999-05-20) the whole document	1-17

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 00 00529

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-10 (partially), 11, 12 (partially)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible. Consequently, the search has been restricted to compounds for which a pharmaceutical use is indicated (particularly as cysteine protease inhibitors).

Claim 11 deals with synthetic intermediates which are amides of di- and tripeptides, a group of compounds which are well-known in the prior art so that it is impossible to determine which parts of the claim may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). No attempt was made to search this claim, but where documents relevant to this claim were found during the search in relation to the other claims, this has been noted in the International Search Report.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Application No
PCT/GB 00/00529

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1073288 A		BE 662788 A CH 456623 A FR 4312 M US 3467691 A US 3697577 A	17-08-1965 16-09-1969 10-10-1972
WO 9924460 A	20-05-1999	AU 1487399 A	31-05-1999

PATENT COOPERATION TREATY

PCT

INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference PHM 70485/WO	FOR FURTHER ACTION		see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.
International application No. PCT/GB 00/ 00529	International filing date (day/month/year) 16/02/2000	(Earliest) Priority Date (day/month/year) 20/02/1999	
Applicant ASTRAZENECA UK LIMITED et al.			

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 6 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

1. Basis of the report

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☒ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

DI- AND TRIPEPTIDE NITRILE DERIVATIVES AS INHIBITORS OF CATHEPSIN L AND CATHEPSIN S

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-10 (partially), 11, 12 (partially)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible. Consequently, the search has been restricted to compounds for which a pharmaceutical use is indicated (particularly as cysteine protease inhibitors).

Claim 11 deals with synthetic intermediates which are amides of di- and tripeptides, a group of compounds which are well-known in the prior art so that it is impossible to determine which parts of the claim may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). No attempt was made to search this claim, but where documents relevant to this claim were found during the search in relation to the other claims, this has been noted in the International Search Report.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

National Application No.

T/GB 00/00529

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D307/68 C07C255/29 C07D333/24 C07K5/062 C07K5/078
 C07K5/065 C07C323/60 C07C317/50 A61K31/275 A61K31/277
 A61K31/341 A61K31/381 A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C C07D C07K A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S. SUZUE, ET AL.: "Hepatic agents. I. Synthesis of aminoacyl (and hydroxyacyl) aminoacetonitriles" CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 16, no. 8, August 1968 (1968-08), pages 1417-1432, XP002108053 Pharmaceutical Society of Japan, Tokyo, JP ISSN: 0009-2363 the whole document <div style="text-align: center;">--- -/--</div>	1-4,6,7, 10-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 June 2000

Date of mailing of the international search report

30/06/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

English, R

INTERNATIONAL SEARCH REPORT

National Application No

T/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A.R. KATRITZKY, ET AL.: "Benzotriazole-assisted synthesis of alpha-aminonitriles and a conceptually novel method for peptide elongation" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 6, June 1990 (1990-06), pages 1853-1857, XP002140042 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X compounds 5g-5m, 6g-6m ---	1,2,4,7, 11
X	GB 1 073 288 A (KYORIN SEIYAKU) 21 June 1967 (1967-06-21) the whole document ---	1,7,10, 12-14
X	S.A. THOMPSON, ET AL.: "Carboxy-modified amino acids and peptides as protease inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 29, no. 1, January 1986 (1986-01), pages 104-111, XP002140043 American Chemical Society, Washington, DC, US ISSN: 0022-2623 compounds 3b, 3d, 4b, 4d ---	1,7, 10-12
X	D.T. ELMORE, ET AL.: "Thioesters of amino acid derivatives as thioacylating agents in thiopeptide synthesis" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 5, May 1988 (1988-05), pages 1051-1055, XP002140044 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X page 1051, right-hand column, line 10 - line 26 ---	1-3,6,7, 10
X	H. MOSER, ET AL.: "Poly(dipeptamidium)-Salze: Definition und Methoden zur präparativen Herstellung" HELVETICA CHIMICA ACTA, vol. 69, 1986, pages 1224-1262, XP002049723 Verlag Helvetica Chimica Aata, Basle, CH ISSN: 0018-019X compounds 3, 4a, 5 --- -/--	1-3,6,7, 10-12

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	D.J. BUTTLE, ET AL.: "Affinity purification of the novel cysteine proteinase papaya proteinase IV, and papain from papaya latex" BIOCHEMICAL JOURNAL, vol. 261, no. 2, 15 July 1989 (1989-07-15), pages 469-476, XP000913659 Portland Press, London, GB ISSN: 0264-6021 page 471, right-hand column ----	1
X	S. LIU, ET AL.: "Effects of ligand homologation and ligand reactivity on the apparent kinetic specificity of papain" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1250, 1995, pages 43-48, XP002140045 Elsevier Science, Amsterdam, NL ISSN: 0006-3002 compound 3a ----	1
X	C.A. LEWIS, ET AL.: "Thiohemiacetal formation by inhibitory aldehydes at the active site" BIOCHEMISTRY, vol. 16, no. 22, 1 November 1977 (1977-11-01), pages 4890-4895, XP002140046 American Chemical Society, Washington, DC, US ISSN: 0006-2960 table II, entries 8, 11 ----	1
P, X	WO 99 24460 A (NOVARTIS) 20 May 1999 (1999-05-20) the whole document -----	1-17


INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/00529

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1073288 A		BE 662788 A CH 456623 A FR 4312 M US 3467691 A US 3697577 A	17-08-1965 16-09-1969 10-10-1972
WO 9924460 A	20-05-1999	AU 1487399 A	31-05-1999



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference DEG/ sjl		FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/GB00/00529	International filing date (day/month/year) 16/02/2000	Priority date (day/month/year) 20/02/1999	
International Patent Classification (IPC) or national classification and IPC C07D307/68			
Applicant ASTRAZENECA AB at al			

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 4 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☐ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 28/08/2000	Date of completion of this report 16.05.2001
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Baston, E Telephone No. +49 89 2399 8229 

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/GB00/00529

I. Basis of the report

1. With regard to the **elements** of the international application (*Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17)*):

Description, pages:

1-63 as originally filed

Claims, No.:

1-17 as originally filed

2. With regard to the **language**, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language: , which is:

- ☐ the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).
☐ the language of publication of the international application (under Rule 48.3(b)).
☐ the language of a translation furnished for the purposes of international preliminary examination (under Rule 55.2 and/or 55.3).

3. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in written form.
☐ filed together with the international application in computer readable form.
☐ furnished subsequently to this Authority in written form.
☐ furnished subsequently to this Authority in computer readable form.
☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/GB00/00529

(Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.)

6. Additional observations, if necessary:

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

☐ the entire international application.

☒ claims Nos. 1-17.

because:

☐ the said international application, or the said claims Nos. relate to the following subject matter which does not require an international preliminary examination (*specify*):

☒ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. 1-17 are so unclear that no meaningful opinion could be formed (*specify*):
see separate sheet

☒ the claims, or said claims Nos. are so inadequately supported by the description that no meaningful opinion could be formed.

☒ no international search report has been established for the said claims Nos. 11 in full; 1-10, 12-17 in part..

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

To section III

The search report states, that claims 1-10 (partially), 11 (in full) and 12 (partially) were not searched.

This report further mentions, that the search for the presently claimed subject-matter revealed such a large number of relevant documents, which necessitated a restriction to those compounds, for which a pharmaceutical use (spec. cysteine protease inhibition) was indicated. Notwithstanding already 10 X-documents were retrieved, which obviously deprive the novelty of the partially searched claims 1-10, 12-14 (compounds) and 15-17 (related methods and use). Thus it is questionable, what is actually the invention (which should be novel and inventive compared with the prior art). Moreover it has to be stated, that the breadth of the claims would render the involvement of an inventive step (Art. 33(3) PCT) a priori doubtful, even if novelty (Art. 33(2) PCT) could be established.

Additionally the IPER wishes to point out, that the drafting of a meaningful opinion over the whole scope of claims, that could only partially be searched is impossible.

Consequently no preliminary opinion is provided for claims 1-17.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DI- AND TRIPEPTIDE NITRILE DERIVATIVES AS INHIBITORS OF CATHEPSIN L AND CATHEPSIN S

The present invention relates to compounds that are cysteine protease inhibitors and in particular compounds that are Cathepsin L inhibitors and or Cathepsin S inhibitors especially
5 Cathepsin S inhibitors. The invention further relates to processes for their preparation, to intermediates useful in their preparation, to their use as therapeutic agents, to pharmaceutical compositions containing them and to a method of treating a Cathepsin L or Cathepsin S mediated disease state.

Cysteine proteases are enzymes important in normal cell physiology, but they are also
10 associated with several disease states including inflammation, metastasis, tissue damage following myocardial infarction, bone resorption and muscle wasting in dystrophic diseases.

Cathepsins B, H, K, L, N and S are cysteinyl proteases involved in normal protein degradation and are normally located in the lysosomes of cells. However, when these enzymes are found outside the lysosomes they have been implicated as playing a causative role in a
15 number of disease states including bone resorption disease such as osteoporosis.

The number of people living to an old age has increased dramatically in recent years. This has been marked by an increase in the number of people having osteoporosis and other diseases associated with old age. Osteoporosis is accompanied by a high incidence of bone fracture resulting in many aged patients being confined to their beds. There is therefore a great
20 need for a pharmaceutical composition to treat or prevent this disease.

Living bone is continuously being remodelled and replenished by the process of resorption and deposition of the protein matrix and calcium minerals. These events are facilitated by the osteoclast, which has the ability to degrade and demineralise the bone, and the osteoblast which is responsible for new bone generation. In normal situations these
25 processes are intimately linked resulting in little alteration of bone mass. However, pathological conditions exist in which there is an imbalance between their activities resulting in increased degradation and demineralisation of bone and the development of fragile and/or brittle bone structure, as seen during osteoporosis. While the exact mechanism for this resorption is not known, increased osteoclast activity, as realised by increased proteolytic
30 activity, is a contributing factor, and selective inhibition of proteolytic action may result in the arrest or reversal of bone loss. The lysosomal cysteine proteinases, Cathepsins B, H, K, L, N


and S have been postulated as the proteinases that are responsible for osteoclast bone resorption, because of their ability to degrade insoluble type I collagens at low pH.

Cathepsins B, H, K, L, N and S have been further implicated as playing a causative role in other diseases such as rheumatoid arthritis, osteoarthritis, tumour metastasis, pneumocystitis, *Crithidia fusiculata*, malaria, *trypanosoma brucei brucei*, schistosomiasis, periodontal disease, metachromatic leukodystrophy and muscular dystrophy. Cathepsins B, H, K, L, N and S, either alone or together, have also been implicated as playing a causative role in chronic obstructive pulmonary disease (COPD).

In recent years a number of synthetic inhibitors of cysteine proteases have been disclosed. US 5,055,451 discloses a series of peptidyl methyl ketones as thiol protease inhibitors; WO 95/15749 discloses peptidyl ketones with heterocyclic leaving groups as cysteine protease inhibitors; the *in vivo* inhibition of Cathepsin B by peptidyl (acyloxy) methyl ketones was discussed in *J. Med. Chem.* **1994**, 37, 1833-40 and these types of compounds as inhibitors of cysteine protease inhibitors were also discussed in *J. Am. Chem. Soc.*, **1988**, 110, 4429-4431; peptidyl diazomethyl ketones as specific inactivators of thiol proteinases was discussed in *J. Biol. Chem.*, **1981**, 256, 4, 1923-8 and in *Methods in Enzymology*, **1981**, 80, 820-5; the inhibiting activities of 1-peptidyl-2-haloacetyl hydrazines towards Cathepsin B and calpains was discussed in *Eur. J. Med. Chem.*, **1993**, 28 297-311 and peptidyl fluoromethyl ketones as inhibitors of Cathepsin B and the implication for treatment of Rheumatoid arthritis was discussed in *Biochemical Pharmacology*, **1992**, 44, 6, 1201-7. Thus, there is a great need for a specific cysteine protease inhibitor, especially a Cathepsin L inhibitor or a Cathepsin S inhibitor.

The present invention discloses compounds with inhibitory activity of cysteine proteases and in particular of Cathepsin L and or Cathepsin S. The compounds of the invention are also useful in the treatment of chronic obstructive pulmonary disease (COPD).

Accordingly the present invention provides a compound of formula (I):



(I)

wherein *r* is 0 or 1;

- R¹** is hydrogen, optionally substituted benzyl where said optional substituents are chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl, or **R¹** is a group of formula (II):




(II)

- wherein **R⁵** is C₁₋₆alkyl (optionally substituted with an optionally substituted phenyl, an optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted phenoxy, optionally substituted phenylsulphonyl, optionally substituted C₃₋₁₂cycloalkyl or Het), C₁₋₆alkoxy, optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted C₃₋₁₂cycloalkyl, Het or optionally substituted phenylC₁₋₆alkoxy; where said optional substituents are chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;
- R²** is H, C₁₋₆alkyl [optionally substituted with one or more of hydroxy, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, R⁴, R⁴C₁₋₆alkylsulphanyl, R⁴C₁₋₆alkylsulphinyl, R⁴C₁₋₆alkylsulphonyl], or **R²** is C₁₋₆alkoxy [optionally substituted with one or more of C₂₋₆alkenyl, C₂₋₆alkynyl, R⁴, R⁴C₂₋₆alkenyl, R⁴C₂₋₆alkynyl, Het and trifluoromethyl], or **R²** is C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxycarbonyl, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, R⁴, R⁴S, R⁴C₁₋₆alkylsulphanyl, *N*-(R⁴C₁₋₆alkyl)carbamoyl, *N*-(HetC₁₋₆alkyl)carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl or C₁₋₆alkylsulphonyl; **R⁴** is an optionally substituted phenyl or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms, said optional

- substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;
- R³** is H or C₁₋₆alkyl;
- (AA¹) and (AA²) are independently chosen from Ala, Arg, Cys, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr, Val,


10




Lys(CHO),

Arg(NO₂),


β-Ala,




Ser(Bzl),




Ph-Gly,



Nle,

Ser(O^tBu),


His(Bzl),




Met(O),

15


5






Cha,




His(Me),




Cit,

Tyr(^tBu),Met(O₂),



Pyr-Ala




Phe(S),




Leu(S),

Phe(CH₂S),


Cy(S)-Gly,




Hetar(S)-Gly,



alk(S)-Gly or




Het(S)-Gly, and

Phe(CH₂CH₂S);

wherein Ring A is C₃₋₁₂cycloalkyl; Ring B is a 5 or 6 membered heteroaryl ring; Ring C is Het; V is C₁₋₆alkyl excluding isopropyl; the nitrogen of the amino acid may optionally be alkylated with C₁₋₆alkyl; the phenyl group of Phe(S) and Rings A and B are optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl or *N,N*-(C₁₋₆alkyl)₂sulphamoyl; the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group; the sulphur moiety in the α -position of the amino acid (AA³) may be optionally oxidised to form an -S(O)₂- or -S(O)- moiety; and Het is a fully saturated monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms;

or a pharmaceutically acceptable salt thereof.

In one aspect the present invention provides a compound of formula (I):




(I)

wherein:

r is 0 or 1;

R¹ is optionally substituted benzyl where said optional substituents are chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl or **R**¹ is a group of formula (II):




(II)

wherein R^5 is C_{1-6} alkyl (optionally substituted with an optionally substituted phenyl, an optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted phenoxy, optionally substituted phenylsulphonyl, optionally substituted C_{3-12} cycloalkyl or Het), C_{1-6} alkoxy, optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted C_{3-12} cycloalkyl, Het, optionally substituted phenyl C_{1-6} alkoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;



R^2 is an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;

R^3 is H; and


(AA^1) and (AA^2) are independently chosen from Ala, Arg, Cys, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr, Val,




Lys(CHO),

Arg(NO₂), β -Ala,


8





Ser(Bzl),




Ph-Gly,




Nle,

Ser(O^tBu),


His(Bzl),




Met(O),






Cha,




His(Me),




Cit,

Tyr(^tBu),Met(O₂),


Pyr-Ala



Phe(S),




Leu(S),


Phe(CH₂S),

5


10




Cy(S)-Gly,



Hetar(S)-Gly,



alk(S)-Gly or



Het(S)-Gly;

- 5 wherein Ring A is C₃₋₁₂cycloalkyl, Ring B is a 5 or 6 membered heteroaryl ring, Ring C is Het, V is C₁₋₆alkyl excluding isopropyl, the nitrogen of the amino acid may optionally be alkylated with C₁₋₆alkyl and the phenyl group of Phe(S) and Rings A and B may be optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino,
- 10 *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl or the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group;
- 15 or a pharmaceutically acceptable salt thereof.

In this specification the term 'alkyl' includes straight chained and branched structures and ring systems. For example, C₁₋₆alkyl includes propyl, isopropyl, *t*-butyl, cyclopropyl and cyclohexyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only, references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only and references to individual cycloalkyl groups such as cyclohexyl are specific to the cyclic groups only.

20

A similar convention applies to other radicals, for example "hydroxyC₁₋₆alkyl" includes 1-hydroxyethyl and 2-hydroxyethyl.

The term "halo" refers to fluoro, chloro, bromo and iodo.

- 25 "Het" means, unless otherwise further specified, a fully saturated monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms. Preferably these ring heteroatoms

are selected from nitrogen, oxygen and sulphur. Examples of "Het" include pyrrolidinyl, imidazolidinyl, pyrazolidinyl, piperidyl, piperazinyl and morpholinyl.

Unless otherwise defined standard amino acid abbreviations are used. For example "Ala" refers to alanine and "Gly" refers to glycine.

5 "5- or 6- membered heteroaryl ring" means, unless otherwise further specified, a 5- or 6- membered ring that contains some degree of unsaturation, with up to four ring heteroatoms selected from nitrogen, oxygen and sulphur. Examples of "5- or 6- membered heteroaryl ring" include thienyl, furyl, imidazolyl, thiazolyl, pyrimidinyl, pyridinyl, pyrrolyl and pyrazolyl. Examples of "6 membered heteroaryl ring" include pyrimidinyl, pyridinyl, pyrazinyl and
10 pyridazinyl. Examples of "5 membered heteroaryl ring" include thienyl, furyl, imidazolyl, thiazolyl, pyrrolyl and oxadiazolyl.

Examples of "C₁₋₆alkanoyloxy" are acetoxy and propionyloxy. Examples of "C₁₋₆alkoxycarbonyl" include methoxycarbonyl, ethoxycarbonyl, *n*- and *t*-butoxycarbonyl. Examples of "C₁₋₆alkoxy" include methoxy, ethoxy and propoxy. Examples of
15 "C₁₋₆alkanoylamino" include formamido, acetamido and propionylamino. Examples of "C₁₋₆alkylsulphanyl" include methylthio and ethylthio. Examples of "C₁₋₆alkylsulphinyl" include methylsulphinyl and ethylsulphinyl. Examples of "C₁₋₆alkylsulphonyl" include mesyl and ethylsulphonyl. Examples of "C₁₋₆alkanoyl" include acetyl and propionyl. Examples of "C₁₋₆alkylamino" include methylamino and ethylamino. Examples of "*N,N*-(C₁₋₆alkyl)₂amino"
20 include *N,N*-dimethylamino, *N,N*-diethylamino and *N*-ethyl-*N*-methylamino. Examples of "*N*-(C₁₋₆alkyl)carbamoylC₁₋₆alkyl" are 2-(methylamino)carbonylethyl and 3-(ethylamino)carbonylpropyl. Examples of "*N,N*-(C₁₋₆alkyl)₂carbamoylC₁₋₆alkyl" are 2-(dimethylamino)carbonylethyl and 3-(*N*-methyl-*N*-ethylamino)carbonylpropyl. Examples of "C₂₋₆alkenyl" are vinyl, allyl and 1-propenyl. Examples of "C₂₋₆alkynyl" are ethynyl,
25 1-propynyl and 2-propynyl. Examples of "*N*-(C₁₋₆alkyl)carbamoyl" are *N*-methylaminocarbonyl and *N*-ethylaminocarbonyl. Examples of "*N,N*-(C₁₋₆alkyl)₂carbamoyl" are *N,N*-dimethylaminocarbonyl and *N*-methyl-*N*-ethylaminocarbonyl. Examples of "*N*-(C₁₋₆alkyl)sulphamoyl" are *N*-methylsulphamoyl and *N*-ethylsulphamoyl. Examples of "*N,N*-(C₁₋₆alkyl)₂sulphamoyl" are *N,N*-dimethylsulphamoyl
30 and *N,N*-diethylsulphamoyl. Examples of "R⁴C₁₋₆alkylsulphanyl" include R⁴methylthio and 2-R⁴ethylthio. Examples of "R⁴C₁₋₆alkylsulphinyl" include R⁴methylsulphinyl and 2-R⁴ethylsulphinyl. Examples of "R⁴C₁₋₆alkylsulphonyl" include R⁴mesyl and 2-

R^4 ethylsulphonyl. Examples of R^4C_{2-6} alkenyl are 2- R^4 vinyl and 3- R^4 allyl. Examples of “ R^4C_{2-6} alkynyl” are 2- R^4 ethynyl and 3- R^4 propyn-1-yl. Examples of “ N -(R^4C_{1-6} alkyl)carbamoyl” are R^4 methylaminocarbonyl and 2- R^4 ethylaminocarbonyl. Examples of “ N -(Het C_{1-6} alkyl)carbamoyl” are morpholinomethylaminocarbonyl and 2-(piperidinoethyl)aminocarbonyl. Examples of “ C_{3-12} cycloalkyl” are cyclopropyl, cyclopentyl and cyclohexyl.

Where optional substituents are chosen from “one or more” groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups. For example where optional substituents are chosen from one or more halo, C_{1-6} alkoxy and C_{1-6} alkyl, examples of possible combinations of substituents include 1) a bromo group, 2) two chloro groups, 3) a methoxy, ethoxy and propoxy substituent, 4) a fluoro and a methoxy group, 5) a methoxy, a methyl and an ethyl group, and 6) a chloro, a methoxy and an ethyl group.

According to a further feature of the invention there is provided a compound of formula (I) wherein:

r is 0 or 1;

R^1 is optionally substituted benzyl where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl or R^1 is a group of formula (II) wherein R^5 is C_{1-6} alkyl (optionally substituted with an optionally substituted phenyl, an optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted phenoxy or optionally substituted phenylsulphonyl), C_{1-6} alkoxy, optionally substituted phenyl, optionally substituted naphthyl, optionally substituted phenyl C_{1-6} alkoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

- R²** is H, C₁₋₆alkyl [optionally substituted with one or more of hydroxy, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, R⁴, R⁴C₁₋₆alkylsulphanyl, R⁴C₁₋₆alkylsulphanyl, R⁴C₁₋₆alkylsulphonyl], or R² is C₁₋₆alkoxy [optionally substituted with one or more of C₂₋₆alkenyl, C₂₋₆alkynyl, R⁴, R⁴C₂₋₆alkenyl, R⁴C₂₋₆alkynyl, Het and trifluoromethyl], or R² is C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxycarbonyl, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, R⁴, R⁴S, R⁴C₁₋₆alkylsulphanyl, *N*-(R⁴C₁₋₆alkyl)carbamoyl, *N*-(HetC₁₋₆alkyl)carbamoyl, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl wherein R⁴ is an optionally substituted phenyl, or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R³ is H or C₁₋₆alkyl; and

- (AA¹) and (AA²) are independently chosen from Ala, Arg, Cys, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr, Val, Lys(CHO), Arg(NO₂), β-Ala, Ser(Bzl), Ph-Gly, Nle, Ser(O^tBu), His(Bzl), Met(O), Cha, His(Me), Cit, Tyr(^tBu), Met(O₂), Pyr-Ala, Phe(S), Leu(S) or Phe(CH₂S); wherein the nitrogen of the amino acid may optionally be alkylated with C₁₋₆alkyl and the phenyl group of Phe(S) may be optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl or the phenyl group may be fused to another phenyl group to form a naphthyl group; or a pharmaceutically acceptable salt thereof.

Preferred values for R¹, r, AA¹, AA², R² and R³ are as follows.

In one aspect of the invention preferably R^1 is optionally substituted benzyl where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl,

5 N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl or R^1 is a group of formula (II) wherein R^5 is C_{1-6} alkoxy, optionally substituted C_{3-12} cycloalkyl or optionally substituted phenyl C_{1-6} alkoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl,
10 halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl.

15 Preferably R^1 is benzyl or a group of formula (II) wherein R^5 is C_{1-6} alkyl (optionally substituted with a 6 membered heteroaryl ring, phenyl, phenylsulphonyl or phenoxy optionally substituted with one or more halo), C_{1-6} alkoxy, phenyl (optionally substituted with one or more halo), naphthyl and phenyl C_{1-6} alkoxy.

In another aspect of the invention preferably R^1 is benzyl or a group of formula (II) wherein R^5 is C_{1-6} alkyl (optionally substituted with a 6 membered heteroaryl ring, phenyl, phenylsulphonyl, C_{3-12} cycloalkyl or phenoxy optionally substituted with one or more halo), C_{1-6} alkoxy, phenyl (optionally substituted with one or more halo), naphthyl, C_{3-12} cycloalkyl, Het, and phenyl C_{1-6} alkoxy.

25 More preferably R^1 is benzyl or a group of formula (II) wherein R^5 is methyl, methoxy, ethoxy, propoxy, ^tbutoxy, phenyl, 2,4-dichlorophenyl, naphthyl, benzyloxy, pyridylmethyl, benzyl, 2,4,6-trichlorophenoxymethyl and phenylsulphonylmethyl.

In another aspect of the invention more preferably R^1 is benzyl or a group of formula (II) wherein R^5 is methyl, methoxy, ethoxy, propoxy, ^tbutoxy, phenyl, 2,4-dichlorophenyl, naphthyl, benzyloxy, pyridylmethyl, benzyl, 2,4,6-trichlorophenoxymethyl,
30 phenylsulphonylmethyl, morpholino, cyclohexyl, cyclopentyl, cyclohexylmethyl and piperidino.

Particularly R^1 is a group of formula (II) wherein R^5 is methyl, 'butoxy, benzyloxy and pyridylmethyl.

In another aspect of the invention particularly R^1 is a group of formula (II) wherein R^5 is methyl, 'butoxy, benzyloxy, pyridylmethyl, morpholino, cyclohexyl, cyclopentyl,
5 cyclohexylmethyl and piperidino.

More particularly R^1 is a group of formula (II) wherein R^5 is methyl, 'butoxy, benzyloxy and 4-pyridylmethyl.

In another aspect of the invention particularly R^1 is a group of formula (II) wherein R^5 is morpholino, cyclohexyl, cyclopentyl, cyclohexylmethyl and piperidino.

10 In one aspect of the invention preferably r is 0.

In another aspect of the invention preferably r is 1.

Preferably AA^1 is Leu, Pyr-Ala and Phe wherein the nitrogen of the amino acid is optionally substituted with C_{1-6} alkyl.

More preferably AA^1 is Leu and the nitrogen of the amino acid is unsubstituted.

15 Preferably AA^2 is Phe, Leu, Ile, Tyr, Tyr('Bu), Val, Cha, Leu(S), Phe(S) and Phe(CH_2S) and the nitrogen of the amino acid is unsubstituted and the phenyl group of Phe(S) is optionally substituted with halo, C_{1-6} alkyl or is fused to another phenyl group to form a naphthyl group.

20 More preferably AA^2 is Tyr, Leu and Phe and the nitrogen of the amino acid is unsubstituted.

Preferred combinations of r , AA^1 and AA^2 are as follows.

When $r = 0$ preferably AA^2 is Phe, Leu, Ile, Val, Tyr, Tyr('Bu), Leu(S), Phe(S) and Phe(CH_2S) and the nitrogen of the amino acid is unsubstituted and the phenyl group of Phe(S) is optionally substituted with halo, C_{1-6} alkyl or is fused to another phenyl group to form a
25 naphthyl group.

When $r = 0$ more preferably AA^2 is Tyr.

When $r = 1$ preferably AA^1-AA^2 is Leu-Leu, Pyr-Ala-Leu, Phe-Leu, Leu-Phe, Leu-Ile, Leu-Val, Leu-Cha and (N-Me)Leu-Leu.

When $r = 1$ more preferably AA^1-AA^2 is Leu-Leu and Leu-Phe.

30 In another aspect of the invention preferably (AA^1) and (AA^2) are both independently selected from Phe(S), Leu(S), Phe(CH_2S), Cy(S)-Gly, Hetar(S)-Gly, alk(S)-Gly and Het(S)-Gly wherein Phe(S) and Rings A and B may be optionally substituted as hereinbefore defined

and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group.

In a further aspect of the invention preferably r is 0 and (AA²) is selected from Phe(S), Leu(S), Phe(CH₂S), Cy(S)-Gly, Hetar(S)-Gly, alk(S)-Gly and Het(S)-Gly wherein Phe(S) and
 5 Rings A and B may be optionally substituted as hereinbefore defined and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group.

In a further aspect of the invention preferably r is 0 and (AA²) is selected from Phe(S), Leu(S) and Phe(CH₂S) wherein Phe(S) may be optionally substituted as hereinbefore defined and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a
 10 naphthyl group.

In one aspect of the invention preferably R² is C₁₋₆alkoxy [optionally substituted with one or more of C₂₋₆alkenyl, C₂₋₆alkynyl, R⁴, R⁴C₂₋₆alkenyl, R⁴C₂₋₆alkynyl, Het and trifluoromethyl], or R² is C₂₋₆alkenyl, C₂₋₆alkynyl, carbamoyl, R⁴, R⁴S, R⁴C₁₋₆alkylsulphanyl, C₁₋₆alkanoylamino, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl wherein R⁴ is
 15 an optionally substituted phenyl, or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl,
 20 *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl.

Preferably R² is hydrogen, C₁₋₆alkyl [optionally substituted with C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl or R⁴], C₁₋₆alkoxy [optionally substituted with C₂₋₆alkynyl] and R⁴-
 25 wherein R⁴ is an optionally substituted phenyl or an optionally substituted 5 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C₁₋₆alkyl and halo.

More preferably R² is hydrogen, methyl, ethyl, propyl, isobutyl, furyl, thienyl, pyrazolyl (optionally substituted with one or more of methyl and bromo), imidazolyl, 1,2,4-triazolyl, phenyl, benzyl, 2-methylthioethyl, methylthio, ethylthio, isopropylthio, mesylethyl, methoxy, ethoxy, isopropoxy and 2-propynyloxy.
 30

Particularly R^2 is furyl, pyrazolyl (optionally substituted with one or more of methyl and bromo), imidazolyl, 1,2,4-triazolyl, benzyl, 2-methylthioethyl, isopropylthio, methoxy, isopropoxy and 2-propynyloxy.

More particularly R^2 is fur-2-yl, pyrazol-1-yl, 3,5-dimethylpyrazol-1-yl, 4-bromo-3,5-dimethylpyrazol-1-yl, imidazol-1-yl, 1,2,4-triazol-1-yl, benzyl, methylthioethyl, isopropylthio, methoxy, isopropoxy and 2-propynyloxy.

In another aspect of the invention preferably R^2 is an optionally substituted 5 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl.

More preferably R^2 is thienyl, furyl and pyrazolyl.

Particularly R^2 is thienyl.

Preferably R^3 is hydrogen.

According to one aspect of the present invention there is provided a compound of the formula (I) wherein:

R^1 is a group of formula (II) wherein R^5 is morpholino, cyclohexyl, cyclopentyl, cyclohexylmethyl and piperidino;

r is 0 or 1;

(AA^1) and (AA^2) are both independently selected from Phe(S), Leu(S), Phe(CH_2S), Cy(S)-Gly, Hetar(S)-Gly, alk(S)-Gly and Het(S)-Gly wherein Phe(S) and Rings A and B may be optionally substituted as hereinbefore defined and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group;

R^2 is an optionally substituted 5 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, N-(C₁₋₆alkyl)sulphamoyl and N,N-(C₁₋₆alkyl)₂sulphamoyl; and

R³ is hydrogen;

or a pharmaceutically acceptable salt thereof.

5 According to another aspect of the present invention there is provided a compound of the formula (I) wherein:

R¹ is benzyl or a group of formula (II) wherein R⁵ is C₁₋₆alkyl (optionally substituted with a 6 membered heteroaryl ring, phenyl, phenylsulphonyl or phenoxy optionally substituted with one or more halo), C₁₋₆alkoxy, phenyl (optionally substituted with one or more halo),
10 naphthyl or phenylC₁₋₆alkoxy;

r is 0 or 1;

AA¹ is Leu, Pyr-Ala or Phe wherein the nitrogen of the amino acid is optionally substituted with C₁₋₆alkyl;

AA² is Phe, Leu, Ile, Tyr, Tyr(^tBu), Val, Cha, Leu(S), Phe(S) and Phe(CH₂S) and the
15 nitrogen of the amino acid is unsubstituted and the phenyl group of Phe(S) is optionally substituted with halo, C₁₋₆alkyl or is fused to another phenyl group to form a naphthyl group;

R² is hydrogen, C₁₋₆alkyl [optionally substituted with C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl or R⁴], C₁₋₆alkoxy [optionally substituted with C₂₋₆alkynyl,] or R⁴ - wherein R⁴ is an optionally substituted phenyl or an optionally substituted 5 membered heteroaryl ring
20 containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C₁₋₆alkyl or halo; and

R³ is hydrogen;

or a pharmaceutically acceptable salt thereof.

A further preferred class of compounds is that of formula (I) wherein:

25 R¹ is a group of formula (II) wherein R⁵ is methyl, ^tbutoxy, benzyloxy or pyridylmethyl;

r is 0 or 1;

AA¹ is Leu wherein the nitrogen of the amino acid is unsubstituted;


AA² is Tyr, Leu or Phe wherein the nitrogen of the amino acid is unsubstituted;

30 R² is furyl, pyrazolyl (optionally substituted with one or more methyl or bromo), imidazolyl, 1,2,4-triazolyl, benzyl, methylthioethyl, isopropylthio, methoxy, isopropoxy or propynyloxy; and

R^3 is hydrogen;

or a pharmaceutically acceptable salt thereof.

In yet another aspect the present invention provides a compound of formula (Ia):




5 wherein:


R^7 is optionally substituted benzyl, optionally substituted phenoxyethyl, optionally substituted phenylsulphonylmethyl, optionally substituted benzyloxy, optionally substituted naphthyl, optionally substituted phenyl or t-butoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;

15 R^6 is hydrogen, optionally substituted phenyl or optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms; said optional substituents being chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;


(AA^3) is selected from:



Phe(S),




Leu(S),




Phe(CH₂S),


19




Cy(S)-Gly,




Hetar(S)-Gly,



alk(S)-Gly,




Het(S)-Gly, and

Phe(CH₂CH₂)S;

- 5 wherein Ring A is C₃₋₁₂cycloalkyl, Ring B is a 5 or 6 membered heteroaryl ring, Ring C is Het, V is C₁₋₆alkyl excluding isopropyl; the nitrogen of the amino acid may optionally be alkylated with C₁₋₆alkyl; the phenyl group of Phe(S) and Rings A and B may be optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino,
- 10 *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl; the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group; and the sulphur moiety in the
- 15 α -position of the amino acid (AA) may be optionally oxidised to form an -S(O)₂- or -S(O)- moiety; and
- Het is a fully saturated monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms;
- or a pharmaceutically acceptable salt thereof.

- 20 In a still further aspect the present invention provides a compound of formula (Ia):




wherein:


R^7 is optionally substituted benzyl, optionally substituted phenoxyethyl, optionally substituted phenylsulphonylmethyl, optionally substituted benzyloxy, optionally substituted naphthyl, optionally substituted phenyl or t-butoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;

R^6 is hydrogen or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;


(AA^3) is selected from:




Phe(S),




Leu(S),




Phe(CH $_2$ S),




Cy(S)-Gly,



Hetar(S)-Gly,



alk(S)-Gly and



Het(S)-Gly;

wherein Ring A is C₃₋₁₂cycloalkyl, Ring B is a 5 or 6 membered heteroaryl ring, Ring C is Het, V is C₁₋₆alkyl excluding isopropyl, the nitrogen of the amino acid may optionally be
 5 alkylated with C₁₋₆alkyl and the phenyl group of Phe(S) and Rings A and B may be optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, N-(C₁₋₆alkyl)carbamoyl, N,N-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto,
 10 C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphonyl, sulphamoyl, N-(C₁₋₆alkyl)sulphamoyl and N,N-(C₁₋₆alkyl)₂sulphamoyl, the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group and the sulphur moiety in the α-position of the amino acid (AA) may be optionally oxidised to form an -S(O)₂- or -S(O)- moiety; or a pharmaceutically acceptable salt thereof. The variable Het is a fully saturated
 15 monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms.

Preferred values for R⁷, AA³ and R⁶ for the compound of formula (Ia) are as follows.

The variable R⁷ is, for example, benzyl (optionally substituted with halo (such as chloro)), α-(C₁₋₄ alkyl)-benzyl (optionally substituted with halo (such as chloro)), α,α-di(C₁₋₄ alkyl)-benzyl (optionally substituted with halo (such as chloro)), optionally substituted
 20 phenoxymethyl, phenylsulphonylmethyl, benzyloxy, naphthyl or optionally substituted phenyl where said optional substituents are chosen from one or more halo.

Preferably R⁷ is benzyl, optionally substituted phenoxymethyl, phenylsulphonylmethyl, benzyloxy, naphthyl or optionally substituted phenyl where said optional substituents are chosen from one or more halo.

25 More preferably R⁷ is benzyl, phenoxymethyl optionally substituted with chloro, phenylsulphonylmethyl, benzyloxy, naphthyl or phenyl optionally substituted with chloro.

Particularly R⁷ is benzyl, 2,4,6-trichlorophenoxymethyl, phenylsulphonylmethyl, benzyloxy, naphthyl or 2,4-dichlorophenyl.

30 More particularly R⁷ is benzyl, 2,4,6-trichlorophenoxymethyl, phenylsulphonylmethyl, benzyloxy, naphth-2-yl or 2,4-dichlorophenyl.

Preferably (AA³) is Leu(S), Phe(S) optionally substituted with C₁₋₆alkyl or halo and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group or the sulphur moiety in the α -position of the amino acid (AA) may be optionally oxidised to form an -S(O)₂- or Phe(CH₂S).

5 More preferably (AA³) is Leu(S), Phe(S), 4-Cl-Phe(S), 3-Cl-Phe(S), 2-Cl-Phe(S), 3-Me-Phe(S), 4-F-Phe(S), Phe(S) fused to another phenyl group to form a naphth-1-yl group, Phe(S) fused to another phenyl group to form a naphth-2-yl group, 4-F-Phe(S) wherein the (S) is oxidised to S(O)₂, 3-Cl-Phe(S) wherein the (S) is oxidised to S(O)₂, 2-Cl-Phe(S) wherein the (S) is oxidised to S(O)₂, 4-Cl-Phe(S) wherein the (S) is oxidised to S(O)₂ or Phe(CH₂S).

10 Preferably R⁶ is hydrogen or a 5 membered heteroaryl ring containing a maximum of four heteroatoms. The heteroatoms are, for example, nitrogen, oxygen or sulphur. The 5 membered heteroaryl ring is, for example, thienyl, furyl, pyrazolyl, imidazolyl or 1,2,4-triazolyl.

Alternatively R⁶ is hydrogen, optionally substituted phenyl or a 5 membered heteroaryl ring containing a maximum of four heteroatoms. The heteroatoms are, for example, nitrogen, oxygen or sulphur. The 5 membered heteroaryl ring is, for example, thienyl, furyl, pyrazolyl, imidazolyl or 1,2,4-triazolyl. Optional substituents include C₁₋₄ alkoxy (such as methoxy).

The variable R⁶ is, for example, methoxyphenyl.

More preferably R⁶ is hydrogen or thienyl.

20 Particularly R⁶ is hydrogen or thien-2-yl.

More particularly R⁶ is thien-2-yl.

In another aspect of the invention preferably R⁶ is an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms wherein said optional substituents are as defined hereinbefore.

25 In another aspect of the invention more preferably R⁶ is an optionally substituted 5 membered heteroaryl ring containing a maximum of four heteroatoms wherein said optional substituents are as defined hereinbefore.

Preferred compounds are those of Examples 1 - 58 or a pharmaceutically acceptable salt thereof.

30 Especially preferred compounds are those of Examples 8, 13, 15, 17, 19, 25, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40 and 41 or a pharmaceutically acceptable salt thereof.

Preferred compounds are those of Examples 59-66 or a pharmaceutically acceptable salt thereof. Especially preferred compounds are those of Examples 59-65.

Preferred compounds are those of Examples 67-74 and 97-113 or a pharmaceutically acceptable salt thereof.

5 Suitable pharmaceutically acceptable salts include acid addition salts such as the methanesulphonate, fumarate, hydrochloride, hydrobromide, citrate and maleate salts and salts formed with phosphoric and sulphuric acid. In another aspect suitable salts are base salts such as an alkali metal salt for example a sodium salt, an alkaline earth metal salt for example a calcium or a magnesium salt, an organic amine salt for example a salt with triethylamine, 10 morpholine, *N*-methylpiperidine, *N*-ethylpiperidine, procaine, dibenzylamine, *N,N*-dibenzylethylamine or an amino acid for example a lysine salt. There may be more than one cation or anion depending on the number of charged functions and the valency of the cations or anions. A preferred pharmaceutically acceptable salt is a sodium salt.

Some compounds of formula (I) may possess chiral centres. It is to be understood that 15 the invention encompasses all such optical isomers and diastereoisomers of compounds of formula (I) which possess cysteine protease inhibitory activity.


The invention further relates to all tautomeric forms of the compounds of formula (I).

It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be 20 understood that the invention encompasses all such solvated forms.

Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt thereof. According to this aspect of the invention there is provided a process (in which variable groups are as defined for formula (I) unless otherwise stated) which comprises:

25


a) coupling an acid of formula (III):



or a reactive derivative thereof;

30 with an amine of formula (IV):

24




(IV)

A suitable reactive derivative of an acid of the formula (III) is, for example, an acyl halide, for example an acyl chloride formed by the reaction of the acid and an inorganic acid chloride, for example thionyl chloride; a mixed anhydride, for example an anhydride formed by the reaction of the acid and a chloroformate such as isobutyl chloroformate; an active ester, for example an ester formed by the reaction of the acid and a phenol such as pentafluorophenol, an ester such as pentafluorophenyl trifluoroacetate, an alcohol such as 1-hydroxybenzotriazole or a uronium salt such as 2-(1-benzotriazolyl)-1,1,3,3-tetramethyluronium hexafluorophosphate(V); an acyl azide, for example an azide formed by the reaction of the acid and an azide such as diphenylphosphoryl azide; an acyl cyanide, for example a cyanide formed by the reaction of an acid and a cyanide such as diethylphosphoryl cyanide; or the product of the reaction of the acid and a carbodiimide such as *N,N*-dicyclohexylcarbodiimide or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

The reaction is preferably carried out in the presence of a suitable base such as, for example, an alkali or alkaline earth metal carbonate, alkoxide or hydroxide, for example sodium carbonate or potassium carbonate, or, for example, an organic amine base such as, for example, pyridine, 2,6-lutidine, collidine, 4-dimethylaminopyridine, triethylamine, morpholine or diazabicyclo-[5.4.0]undec-7-ene. The reaction is also preferably carried out in a suitable inert solvent or diluent, for example methylene chloride, acetonitrile, tetrahydrofuran, 1,2-dimethoxyethane, *N,N*-dimethylformamide, *N,N*-dimethylacetamide, *N*-methylpyrrolidin-2-one or dimethylsulphoxide, and at a temperature in the range, for example, -78° to 150°C, conveniently at or near ambient temperature.

b) dehydrating a compound of formula (V):




(V)

under standard conditions.

For example such a dehydration reaction may conventionally be carried out by reaction with a reagent such as trifluoroacetic anhydride. The reaction can conveniently be conducted in the presence of a suitable base as defined hereinbefore such as, for example, triethylamine. The reaction is also preferably carried out in a suitable inert solvent or diluent, as defined hereinbefore such as dichloromethane and at a temperature in the range, for example, -10°C to reflux conveniently 10°C to reflux.


c) for compounds of formula (I) where $r = 1$, coupling an acid of formula (VI):



(VI)

or a reactive derivative thereof as defined hereinbefore;


with an amine of formula (VII):



(VII)


The reaction can conveniently be conducted under standard coupling conditions, such as those described in a) above.

d) For compounds of formula (I) where R^1 is a group of formula (II) reaction of an amine of formula (VIII):



(VIII)

with an acid of formula (IX):




(IX)


or a reactive derivative thereof as defined hereinbefore.

The reaction can conveniently be conducted under standard coupling conditions, such as those described in a) above.

- 5 e) Compounds of formula (I) where R¹ is optionally substituted benzyl may be obtained by reaction of an amine of formula (X):



- i) with a compound of formula (XI):




where (R)_n are optional substituents as defined above and L is a displaceable group.

A suitable displaceable group L is, for example, a halogeno or sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or toluene-4-sulphonyloxy group.

- 15 This reaction may be carried out under standard conditions such as, for example, those described in *Synthesis* **1993**, 12, 1243-6; or

- ii) by reaction with an aldehyde of formula (XII):




- 20 This reaction may be carried out under standard conditions such as, for example, those described in *Synth. Commun.*, **1995**, 25, 18, 2819-2827.

- Another aspect of the present invention provides a process for preparing a compound of formula (Ia) or a pharmaceutically acceptable salt thereof. According to this aspect of the invention there is provided a process (in which variable groups are as defined for formula (Ia) unless otherwise stated) which comprises:

- a) coupling an acid of formula (IIIa):


27



(IIIa)


or a reactive derivative thereof;

with an amine of formula (IVa):



(IVa)


b) dehydrating a compound of formula (Va):



(Va)


10 under standard conditions; or,

c) reaction of an amine of formula (VIa):



(VIa)

with an acid of formula (VIIa):



(VIIa)

or a reactive derivative thereof (as defined hereinbefore). These processes can be conducted as hereinbefore described.

If not commercially available, the necessary starting materials for the procedures described above may be made by procedures which are selected from standard organic chemical techniques, techniques which are analogous to the synthesis of known, structurally similar compounds, by techniques which are analogous to the above described procedures or by techniques which are analogous to the procedures described in the examples.

For example, it will be appreciated that certain of the optional substituents on a phenyl or naphthyl or a heteroaryl ring in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and a Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications include the reduction of a nitro group to an amino group by, for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an

arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

Many of the intermediates defined herein are novel, for example, those of the formula (V) and these are provided as a further feature of the invention. Moreover some of the starting materials for use in process variant (b) described hereinbefore, namely those compounds of the formula (VIII) are not only novel but also active as inhibitors of Cathepsin L and or Cathepsin S. Accordingly these compounds are provided as a further feature of the invention.

According to a further feature of the invention there is provided a compound of the formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, for use in a method of treatment of the human or animal body by therapy.

In order to use a compound of the formula (I) or (Ia) or a pharmaceutically acceptable salt thereof for the therapeutic treatment of mammals including humans, in particular in the

inhibition of a cysteine protease, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

In another aspect the present invention provides a pharmaceutical composition comprising a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof,
5 and a pharmaceutically acceptable diluent or carrier.

In a further aspect the present invention provides a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, for use as a medicament.

In a still further aspect the present invention provides the use of a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, in the manufacture of a
10 medicament for use in the inhibition of a cysteine protease in a warm blooded animal, such as man.

In a still further aspect the present invention provides the use of a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of chronic obstructive pulmonary disease in a warm
15 blooded animal, such as man.

A method of treating a Cathepsin L or Cathepsin S mediated disease state in mammals which comprises administering to a mammal in need of such treatment an effective amount of a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof.

According to a further feature of the present invention there is provided a method for
20 producing inhibition of a cysteine protease in a warm blooded animal, such as man, in need of such treatment, which comprises administering to said animal an effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof.

In particular the invention provides the use of a compound of the formula (I) or (Ia) of the present invention, or a pharmaceutically acceptable salt thereof, in the manufacture of a
25 medicament for use in the inhibition of Cathepsin S in a warm blooded animal, such as man.

In order to use a compound of the formula (I) or (Ia) or a pharmaceutically acceptable salt thereof for the therapeutic treatment of mammals including humans, in particular in the inhibition of a cysteine protease, it is normally formulated in accordance with standard pharmaceutical practice as a pharmaceutical composition.

30 The pharmaceutical compositions of this invention may be administered in standard manner for the disease condition that it is desired to treat, for example by oral, rectal or parenteral administration. For these purposes the compounds of this invention may be

formulated by means known in the art into the form of, for example, tablets, capsules, aqueous or oily solutions or suspensions, (lipid) emulsions, dispersible powders, suppositories, ointments, creams, drops and sterile injectable aqueous or oily solutions or suspensions.

5 A suitable pharmaceutical composition of this invention is one suitable for oral administration in unit dosage form, for example a tablet or capsule which contains between 100 mg and 1 g of the compound of this invention.

In another aspect a pharmaceutical composition of the invention is one suitable for intravenous, subcutaneous or intramuscular injection.

10 Each patient may receive, for example, an intravenous, subcutaneous or intramuscular dose of 1 mgkg^{-1} to 100 mgkg^{-1} of the compound, preferably in the range of 5 mgkg^{-1} to 20 mgkg^{-1} of this invention, the composition being administered 1 to 4 times per day. The intravenous, subcutaneous and intramuscular dose may be given by means of a bolus injection. Alternatively the intravenous dose may be given by continuous infusion over a period of time. Alternatively each patient will receive a daily oral dose which is approximately
15 equivalent to the daily parenteral dose, the composition being administered 1 to 4 times per day.

The following illustrate representative pharmaceutical dosage forms containing the compound of formula (I) or (Ia), or a pharmaceutically-acceptable salt thereof (hereafter compound X), for therapeutic or prophylactic use in humans:

20 (a)

<u>Tablet I</u>	<u>mg/tablet</u>
Compound X.	100
Lactose Ph.Eur.	179
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

(b)

<u>Tablet II</u>	<u>mg/tablet</u>
Compound X	50
Lactose Ph.Eur.	229
Croscarmellose sodium	12.0
Polyvinylpyrrolidone	6
Magnesium stearate	3.0

(c)

<u>Tablet III</u>	<u>mg/tablet</u>
Compound X	1.0
Lactose Ph.Eur.	92
Croscarmellose sodium	4.0
Polyvinylpyrrolidone	2.0
Magnesium stearate	1.0

5 (d)

<u>Capsule</u>	<u>mg/capsule</u>
Compound X	10
Lactose Ph.Eur.	389
Croscarmellose sodium	100
Magnesium stearate	1.

(e)

<u>Injection I</u>	<u>(50 mg/ml)</u>
Compound X	5.0% w/v
Isotonic aqueous solution	to 100%

10 Buffers, pharmaceutically-acceptable cosolvents such as polyethylene glycol, polypropylene glycol, glycerol or ethanol or complexing agents such as hydroxy-propyl β cyclodextrin may be used to aid formulation.

Note

The above formulations may be obtained by conventional procedures well known in the pharmaceutical art. The tablets (a)-(c) may be enteric coated by conventional means, for example to provide a coating of cellulose acetate phthalate.

5

Inhibition of Cathepsin L and S.

The pharmaceutically-acceptable compounds of the present invention are useful in the inhibition of Cathepsin L and Cathepsin S, having a good activity *in vitro* against human Cathepsin L, human Cathepsin S and rabbit Cathepsin L.

10

Cathepsin L Assay

Recombinant human Cathepsin L was cloned and expressed in E Coli and purified using the method as described by Zeneca Limited, GB 2 306 961 A (published 14.05.1997).

Rabbit Cathepsin L was purified from rabbit liver as described by Maciewicz R. A. and Etherington D. J. (Biochem. J. (1988) 256, 433-440) except the liver homogenate supernatant was concentrated by fractionation with $(\text{NH}_4)_2\text{SO}_4$ (20-80% saturation), and the pellet taken up and dialysed against 20mM NaAcetate pH 5.5, 1mM ethylenediaminetetraacetic acid (EDTA). The supernatant was then applied to a CM Sepharose ion exchange column and Cathepsin L eluted by gradient elution (0.25-0.75M NaCl). Fraction activity was determined using the synthetic substrate NCBz-Phe-Arg-NHMec as described. Cathepsin L fractions were pooled and desalted on a Sephacryl S100 column. Active fractions were pooled, adjusted to 20% saturation $(\text{NH}_4)_2\text{SO}_4$ and concentrated on a phenyl sepharose column. The remaining purification steps were as described.

Cathepsin L activity was measured based on the method of Barrett and Kirschke (1981 Methods in Enzymology, **80**, 535-561), using the fluorogenic substrates NCBz-Phe-Arg-NHMec. Inhibitors were identified by their ability to decrease the generation of the fluorescent leaving group (NHMec). Briefly the assay was as follows:

rHuman Cathepsin L or rabbit Cathepsin L (0.025 pmoles) was pre-incubated with or without test compound in 0.1M sodium acetate buffer pH4.5, 10mM cysteine, 0.1% Brij 35 at 25°C for 15 minutes in a solid black 96 well plate. Synthetic substrate, 20µM NCBz-Phe-Arg-NHMec, was added and the mixture incubated at 37°C for 30 minutes. The reaction was stopped by the addition of 0.1M sodium chloroacetate pH 4.3. Fluorescence was determined

30

using a Fluoroskan II plate reader; excitation 355nm, emission 460nm. Compound potency was determined from the raw fluorescence data by calculating the IC₅₀ against each enzyme using a PC graph drawing software package.

5 Cathepsin S assay.

Cloning and Expression of human Cathepsin S.

Recombinant human Cathepsin S was cloned and expressed in Baculovirus, by the following method. The cDNA sequence for human Cathepsin S is available in the EMBL database Accession Number M90696. This database sequence was used to prepare, by PCR
10 on mRNA from human tissues, a recombinant plasmid carrying an insert with a DNA sequence identical to that of Cathepsin S in the EMBL database (Acc No M90696). The techniques for mRNA isolation, PCR and cloning are standard techniques known by those skilled in the art. Sequence determination of the recombinant insert was carried out using established DNA sequencing techniques.

15 The PCR was done so as to introduce an EcoRI cloning site 5' of the 'ATG' of Cathepsin S and an XbaI cloning site 3' of the 'Stop' codon. The PCR product was cloned between the EcoRI and XbaI sites of the baculovirus transfer vector pFASTBAC-1 (Bac-to-Bac Expression System commercially available from Gibco BRL –Life Technologies (cat no 10359-016)). This recombinant construct was used to generate, by standard techniques,
20 recombinant baculovirus capable of expressing preprocathepsin S.

Expression of recombinant Cathepsin S was tested for the baculoviral constructs by infection of two insect cell lines : Sf9 cells (ATCC No CRL-1711) and T.ni cells (Invitrogen, Cat No B855-02).

25 Purification of Cathepsin S

Method 1.

Procathepsin S was found in the insect cell medium and acid activated. The medium was mixed with an equal volume of 100mM Sodium Acetate buffer pH 4.5, 5mM dithiothreitol (DTT) and 5mM EDTA and incubated for one hour at 37°C method of Maubach
30 et al (Eur. J. Biochem., 250, 745-750, 1997).

Method 2.

The pH of insect cell medium (10ml) containing procathepsin S was adjusted to 4.5 with glacial acetic acid and DTT and EDTA added to 5mM. The sample was then incubated at 37°C for 150min to enable conversion to the active enzyme. Ammonium sulphate was then added to 80% saturation and a pellet obtained by centrifugation. This pellet was redissolved in 2ml buffer A (100mM Tris, 500mM NaCl, 1mM EDTA, pH7.5) and mixed in a batchwise fashion with 100µl thiopropyl-Sepharose for 15min at 4°C. The non bound fraction was removed by a brief centrifugation and the gel washed with 2x1ml buffer A. Cathepsin S was then eluted by batch mixing with 0.4ml 20mM DTT in buffer A for 15min at 4°C.

Measurement of Cathepsin S Activity.

Cathepsin S activity was measured based on the method of Maubach et al (Eur. J. Biochem., 250, 745-750, 1997), using the fluorogenic substrate Z-Val-Val-Arg-NHMec. Inhibitors were identified by their ability to decrease the generation of the fluorescent leaving group (NHMec). Briefly the assay was as follows:

rHuman Cathepsin S (1.5 nmoles) was pre-incubated with or without compounds in 50mM Potassium phosphate buffer pH 6.0-6.2, 20mM Na₂EDTA, 0.1% Brij at 25°C for 5 minutes in a solid black 96 well plate. Synthetic substrate, 20µM Z-Val-Val-Arg-NHMec, was added and the mixture incubated at 30°C for 20 minutes. The reaction was stopped by the addition of 0.1M sodium chloroacetate pH 4.3. Fluorescence was determined using a Fluoroskan II plate reader; excitation 355nm, emission 460nm. Compound potency was determined from the raw fluorescence data by calculating the IC₅₀ against Cathepsin S using a PC graph drawing software package.

The following results were obtained on a standard *in-vitro* test system for the inhibition of Cathepsin L. The activity is described in terms of IC₅₀.

When tested in the above *in-vitro* tests the compounds of this invention give IC₅₀s in the range 1-10,000 nM.

The following data was obtained for Examples 1, 19 and 26:

Example	IC ₅₀ (Human) (nM)	IC ₅₀ (Rabbit) (nM)
1		297
19	38	38.27
26	5651	

The invention will now be illustrated by the following non-limiting examples in which, unless stated otherwise:

- 5 (i) temperatures are given in degrees Celsius (°C); operations were carried out at room or ambient temperature, that is, at a temperature in the range of 18-25°C;
- (ii) organic solutions were dried over anhydrous magnesium sulphate; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (600-4000 Pascals; 4.5-30 mm Hg) with a bath temperature of up to 60°C;
- 10 (iii) chromatography means flash chromatography on silica gel; thin layer chromatography (TLC) was carried out on silica gel plates; where a "Bond Elut" column is referred to, this means a column containing 10g or 20g of silica of 40 micron particle size, the silica being contained in a 60ml disposable syringe and supported by a porous disc, obtained from Varian, Harbor City, California, USA under the name "Mega Bond Elut SP";
- 15 (iv) in general, the course of reactions was followed by TLC and reaction times are given for illustration only;
- (v) final products had satisfactory proton nuclear magnetic resonance (NMR) spectra;
- (vi) yields are given for illustration only and are not necessarily those which can be obtained by diligent process development; preparations were repeated if more material was required;
- 20 (vii) when given, NMR data is in the form of delta values for major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard, determined at 250 MHz using perdeuterio dimethyl sulphoxide (DMSO- δ_6) as the solvent unless otherwise stated;
- (viii) chemical symbols have their usual meanings; SI units and symbols are used;
- 25 (ix) solvent ratios are given in percentage by volume;
- (x) mass spectra (MS) were run with an electron energy of 70 electron volts in the chemical ionisation (CI) mode using a direct exposure probe; where indicated ionisation was effected

by electron impact (EI) or fast atom bombardment (FAB); where values for m/z are given, generally only ions which indicate the parent mass are reported;

(xi) melting points are uncorrected and (dec) indicates decomposition; the melting points given are those obtained for the materials prepared as described; polymorphism may result in

5 isolation of materials with different melting points in some preparations; and


(xii) Z refers to benzyloxycarbonyl and Boc refers to *tert*-butoxycarbonyl.

Example 1


2-(Boc-L-phenylalanyl)-2-(2-furyl)acetonitrile









10 A mixture of Boc-L-phenylalanine (5.0 g), 2-(2-furyl)acetonitrile (3.0 g), hydroxybenzotriazole (5.1 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (4.0 g) and triethylamine (2.63 ml) in *N,N*-dimethylformamide (950 ml) was stirred at 0°C for 30 minutes and then at ambient temperature for 14 hours. The solvent was removed under reduced pressure and the residue was partitioned between ethyl acetate (100 ml) and water
15 (100 ml). The ethyl acetate layer was separated and washed with 2M hydrochloric acid, brine, saturated aqueous sodium bicarbonate and dried. The residue obtained on removal of the solvent was subjected to chromatography on silica by elution with a mixture of dichloromethane and ethyl acetate (9:1 v/v) to give 2-(Boc-L-phenylalanyl)-2-(2-furyl)acetonitrile (4.53 g). Mp 157-158°C; m/z 370 (MH^+); NMR ($CDCl_3$) 1.4 (s, 9H), 3.1 (m,
20 2H), 4.37 (q, 1H), 4.87 (m, 1H), 6.1 (d, 1H), 6.39 (m, 1H), 6.45 (d, 1H), 6.72 (m, 1H), 7.22 (m, 5H), 7.41 (d, 1H).

Using this method but with appropriate starting materials there were prepared the following compounds:



25

Example	R^1	AA^2	R^2	MH^+	Mp (°C)
2	^t BuO-	Leu		336	

3	Me	Leu			152-4
4	Me	Leu			95-8
5	^t BuO-	Ile			106-8
6	^t BuO-	Val			106-9
7	^t BuO-	Leu	H		110-2
8	PhCH ₂ O-	Tyr		420	
9	PhCH ₂ O-	Tyr(^t Bu)			97-100
10	PhCH ₂ O-	Tyr			140
11	PhCH ₂ O-	Phe			131-2
12	^t BuO-	Leu	Ph	346	120-121


Example 13**(2S)-2-(Z-Leu-Leu-NH)-3-phenylpropionitrile**

Trifluoroacetic anhydride (0.28 ml) was added dropwise to a mixture of Z-Leu-Leu-Phe-NH₂ (0.9 g) and pyridine (10 ml) which was stirred under argon at -10°C. The mixture was allowed to warm to room temperature over 1 hour, diluted with water and extracted with ethyl acetate. The extract was washed successively with 1M hydrochloric acid and brine, dried and evaporated to dryness and the residue was recrystallized from ethyl acetate/hexane to give (2S)-2-(Z-Leu-Leu-NH)-3-phenylpropionitrile (0.57 g). Mp 152-154°C; m/z 507 (MH)⁺;

NMR 0.75-0.95 (m, 12H), 1.3-1.7 (m, 6H), 3.05 (d, 2H), 4.05 (m, 1H), 4.3 (m, 1H), 4.9 (m, 1H), 7.2-7.45 (m, 11H), 7.85 (d, 1H), 8.75 (d, 1H).

Example 14-19:

- 5 The following compounds were prepared by a similar process to that described in Example 13:



Example	R ¹	AA ¹	AA ²	R ²	Mp (°C)	MH ⁺
14 ^{1,2}	BOC	Leu	Leu	MeSCH ₂ CH ₂ -	129-130	457
15 ^{1,2}	Z	Leu	Leu	MeSCH ₂ CH ₂ -	-	491
16 ^{1,2}	PhCO	Pyr-Ala	Leu	MeSCH ₂ CH ₂ -	-	496
17 ^{1,3}	Z	Leu	Leu	i-PrS-	-	491
18 ^{1,3}	Z	Phe	Leu	i-PrS-	-	525
19 ^{1,3}	Z	Leu	Phe	i-PrS-	163-165	525

Example 20

- 10 (2S)-2-(Z-Phe-Leu-NH)-4-methylthiobutyronitrile

Phosphoryl chloride (0.07 ml) was added dropwise to stirred, ice-cooled *N,N*-dimethylformamide (2 ml) and the resulting solution added to a stirred ice cooled solution of Z-Phe-Leu-Met-NH₂ (163 mg) in *N,N*-dimethylformamide (2 ml) under an atmosphere of argon. The mixture was stirred for 0.5 hours then poured into ice-water and extracted with ethyl acetate. The extract was washed with water, dried and evaporated to dryness. The residue was recrystallized from ethyl acetate and hexane to give (2S)-2-(Z-Phe-Leu-NH)-4-methylthiobutyronitrile (64 mg). Mp 140-144°C; m/z 525 (MH)⁺; NMR 0.7-0.95 (m, 6H), 1.1-1.5 (m, 3H), 2.0-2.2 (m, 5H), 2.45-2.65 (m, 2H), 2.7-3.0 (m, 2H), 4.25 (m, 2H), 4.65 (q, 1H), 5.0 (m, 2H), 7.1-7.45 (m, 10H), 7.7 (d, 1H), 8.3 (d, 1H), 8.5 (d, 1H).

¹ Purified by flash chromatography on silica (Merck, ART 9385) using mixtures of ethyl acetate and hexane as eluent.

² S isomer at aminoacetonitrile.

³ mixture of epimers at aminoacetonitrile

Example 21**(2S)-2-[(N-phenylacetyl-N-methyl-Leu)-Leu-NH]-4-methylthiobutyronitrile**

The process described in Example 20 was repeated using (2S)-2-[(N-phenylacetyl-N-methyl-Leu)-Leu-Met-NH₂] instead of Z-Phe-Leu-Met-NH₂ as starting material to give (2S)-2-
5 [(N-phenylacetyl-N-methyl-Leu)-Leu-NH]-4-methylthiobutyronitrile. M/z 489 (MH)⁺; NMR [at 100 °C] 0.9 (m, 12H), 1.4-1.8 (m, 6H), 2.0-2.2 (m, 5H), 2.55 (m, 2H), 2.95 (m, 3H), 3.75 (s, 2H), 4.35 (m, 1H), 4.75-5.0 (m, 2H), 7.1-7.5 (m, 6H), 8.25 (m, 1H).

Example 22**(2S)-2-[(N-benzoyl-N-methyl-Leu)-Leu-NH]-4-methylthiobutyronitrile**

The process described in Example 20 was repeated using (2S)-2-[(N-benzoyl-N-methyl-Leu)-Leu-Met-NH₂] instead of Z-Phe-Leu-Met-NH₂ as starting material to give (2S)-2-
10 [(N-benzoyl-N-methyl-Leu)-Leu-NH]-4-methylthiobutyronitrile. M/z 475 (MH)⁺; NMR [at 100 °C] 0.9 (m, 12H), 1.5-1.8 (m, 6H), 2.0-2.2 (m, 5H), 2.55 (m, 2H), 2.9 (m, 3H), 4.4 (m, 1H), 4.75-5.0 (m, 2H), 7.3-7.6 (m, 6H), 8.3 (m, 1H).

Example 23**(2S)-2-(Z-Leu-Leu-NH)-4-methylsulphonylbutyronitrile**

A mixture of (2S)-2-(Z-Leu-Leu-NH)-4-methylthiobutyronitrile (80 mg), oxone (150
20 mg), ethanol (2 ml) and water (1 ml) was stirred at room temperature for 18 hours. The mixture was diluted with water and extracted with ethyl acetate. The extract was dried and evaporated to dryness. The residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent to give (2S)-2-(Z-Leu-Leu-NH)-4-methylsulphonylbutyronitrile (80 mg). M/z 489 (MH)⁺; NMR
25 (CDCl₃) 0.8-1.0 (m, 12H), 1.3-1.6 (m, 15H), 2.4 (m, 2H), 3.0 (s, 3H), 3.2 (m, 2H), 4.05 (m, 1H), 4.35 (m, 1H), 4.7-5.05 (m, 2H), 6.55 (d, 1H), 7.7 (d, 1H).

Example 24**(2RS)-2-(Z-Leu-Leu-NH)-2-phenylacetoneitrile**


30 A mixture of (2RS)-2-(BOC-LeuNH)-2-phenylacetoneitrile (0.69 g), dichloromethane (10 ml) and N,N-diisopropylethylamine (0.5 ml) was stirred under an argon atmosphere and iodotrimethylsilane (0.36 g) was added dropwise. The mixture was stirred for 1 hour and then

additional iodotrimethylsilane (0.36 g) was added and the mixture was stirred for a further hour. N-methylmorpholine (0.5 ml) was added followed by methanol (0.5 ml) and then the solution was evaporated to dryness.

A mixture of the residue, *N,N*-dimethylformamide (10 ml), Z-LeuOH (0.58 g), 1-hydroxybenzotriazole (0.3 g), 1-dimethylaminopropyl-3-ethylcarbodiimide (0.42 g) and N-methylmorpholine (1 ml) was stirred at room temperature for 18 hours. The mixture was evaporated to dryness and the residue dissolved in ethyl acetate, and the solution was washed successively with 0.5M hydrochloric acid, 1M sodium hydroxide and brine and then dried and evaporated to dryness. The residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent to give (2*RS*)-2-(Z-Leu-Leu-NH)-phenylacetonitrile (0.45 g). Mp 168-170°C; *m/z* 493 (MH)⁺; NMR 0.75-0.95 (m, 12H), 1.25-1.75 (m, 6H), 4.05 (m, 1H), 4.4 (m, 1H), 5.0 (m, 2H), 6.15 (m, 1H), 7.3-7.5 (m, 11H), 7.95 (m, 1H), 9.15-9.3 (m, 1H).

15 Examples 25-29:

The following compounds were prepared by a similar process to that described in Example 24:



Example	R ¹	AA ¹	AA ²	R ²	MH ⁺
25	Boc	Leu	Leu	furan-2-yl	449
26	Boc	Leu	Phe	furan-2-yl	483
27	Boc	Leu	Ile	furan-2-yl	449
28	Boc	Leu	Val	furan-2-yl	435
29	Z	Leu	Leu	H	417

20 Example 30


(2*RS*)-2-(Z-Leu-Leu-NH)-2-methoxyacetonitrile

N-bromosuccinimide (133 mg) was added to a stirred solution of (2*RS*)-2-(Z-Leu-Leu-NH)-2-(2-propylthio)acetonitrile (245 mg) in methanol (10 ml), the mixture was allowed to

warm to room temperature and it was then stirred at room temperature for 1 hour. The mixture was evaporated to dryness and the residue was partitioned between ethyl acetate and water. The ethyl acetate was dried and evaporated to dryness and the residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent and the product was recrystallized from carbon tetrachloride to give (2RS)-2-(Z-Leu-Leu-NH)-2-methoxyacetonitrile (190 mg). M/z 447 (MH)⁺; NMR 0.75-0.95 (m, 12H), 1.25-1.75 (m, 6H), 3.3 (s, 3H), 4.05 (m, 1H), 4.2-4.4 (m, 1H), 5.0 (s, 2H), 5.95 (m, 1H), 7.3-7.4 (m, 6H), 7.95 (m, 1H), 9.5-9.7 (m, 1H).

10 Example 31-33:

The following compounds were prepared by a similar process to that described in Example 30:



Example	R ¹	AA ¹	AA ²	R ²	Mp (°C)	MH ⁺
31 ⁴	Z	Leu	Leu	i-PrO		475
32 ^{4,5}	Z	Leu	Leu	2-propynyloxy		471
33 ^{4,5}	Z	Leu	Leu	Pyrazol-1-yl	183-184	483

15 Example 34

(2RS)-2-(Z-Leu-Phe-NH)-2-(pyrazol-1-yl)acetonitrile

A mixture of (2RS)-2-(Z-Leu-Phe-NH)-2-(2-propylthio)acetonitrile (105 mg), yellow mercuric oxide (100 mg), pyrazole and tetrahydrofuran (5 ml) was stirred at room temperature for 18 hours. The mixture was filtered and the filtrate was dried and evaporated to dryness.


20 The residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent and the product was recrystallized from a mixture of dichloromethane and hexane to give (2RS)-2-(Z-Leu-Phe-NH)-2-(pyrazol-1-yl)acetonitrile (70 mg). Mp 189-192 °C; m/z 517 (MH)⁺; NMR 0.75-0.95 (m, 6H), 1.2-1.7 (m, 3H), 2.7-3.1 (m, 2H), 4.0 (m, 1H), 4.6 (m, 1H), 5.05 (s, 2H), 6.35 (m,

⁴ mixture of epimers at aminoacetonitrile (NB varying ratios of isomers)

1H), 7.0-7.5 (m, 12H), 7.65 (m, 1H), 7.7-7.85 (m, 1H), 7.95-8.1 (m, 1H), 10.05-10.25 (m, 1H).

Examples 35-38:

- 5 The following compounds were prepared by a similar process to that described in Example 34:



Example	R ¹	AA ¹	AA ²	R ²	MH ⁺
35	Z	Leu	Leu	imidazol-1-yl	483
36	Z	Leu	Leu	1,2,4-triazol-1-yl	484
37	Z	Leu	Leu	3,5-dimethyl-pyrazol-1-yl	511
38	Z	Leu	Leu	4-bromo-3,5-dimethyl-pyrazol-1-yl	589

Example 39

10 2-(Z-Leu-Cha-NH)acetonitrile

- A mixture of Z-Leu-Cha-OH (0.95 g), aminoacetonitrile hydrochloride (0.23 g), 1-hydroxybenzotriazole (0.46 g), 1-dimethylaminopropyl-3-ethylcarbodiimide (0.5 g), *N,N*-dimethylformamide (6 ml) and *N*-methylmorpholine (0.75 ml) was stirred at room temperature for 18 hours. The mixture was diluted with ethyl acetate, and the solution was washed successively with 0.5 M hydrochloric acid, 1M sodium hydroxide and brine and then dried and evaporated to dryness. The residue was purified by flash chromatography on silica (Merck, ART 9385) using a 49:1 v:v mixture of dichloromethane and methanol as eluent and fractions containing the required product were collected, dried and evaporated to dryness. The residue was triturated with ether and the insoluble white solid collected to give 2-(Z-Leu-Cha-NH)acetonitrile (0.46 g). *M/z* 493 (MH)⁺; NMR 0.75-0.95 (m, 6H), 1.0-1.8 (m, 14H), 4.0-4.2 (m, 3H), 4.35 (q, 1H), 5.05 (s, 2H), 7.25-7.5 (m, 6H), 7.95 (d, 1H), 8.6 (m, 1H).
- 15
- 20

⁵ 5 equivalents of alcohol/pyrazole in tetrahydrofuran solution and 18 hours reaction time.

Example 40**2-[[2-(4-Pyridyl)acetyl]-Leu-Leu]-4-methylthiobutyronitrile**

A mixture of 2-(H-Leu-Leu-NH)-4-methylthiobutyronitrile (71 mg), 1-hydroxybenzotriazole (34 mg), 1-dimethylaminopropyl-3-ethylcarbodiimide (48 mg), 2-(4-pyridyl)acetic acid hydrochloride (35 mg), *N,N*-dimethylformamide (2 ml) and *N*-methylmorpholine (0.2 ml) was stirred at room temperature for 18 hours. The mixture was diluted with ethyl acetate, and the solution was washed successively with 1M sodium hydroxide and brine, dried and evaporated to dryness. The residue was recrystallized from ethyl acetate to give 2-[[2-(4-pyridyl)acetyl]-Leu-Leu]-4-methylthiobutyronitrile (42 mg). Mp 177-178 °C; M/z 476 (MH)⁺; NMR 0.75-0.95 (m, 12H), 1.3-1.7 (m, 6H), 1.9-2.1 (m, 5H), 2.5 (m, 2H), 3.5 (s, 2H), 4.15-4.4 (m, 2H), 4.85 (q, 1H), 7.25 (d, 2H), 8.1 (d, 1H), 8.35 (d, 1H), 8.45 (d, 2H), 8.65 (d, 1H).

Example 41**Ac-Leu-Leu-(2-furyl)acetonitrile**

Trimethylsilyl iodide (0.7 ml) was added to a solution of Boc-Leu-Leu-(2-furyl)-acetonitrile (1.68 g) in chloroform (50 ml) at 0°C. The mixture was stirred at 0°C for 15 minutes and the solvent was removed under reduced pressure. The residue was dissolved in pyridine (20 ml), the solution was cooled to 0°C and acetic anhydride (20 ml) was added and the reaction mixture was stirred at ambient temperature for 14 hours. The reaction mixture was evaporated to dryness and the residue was dissolved in ethyl acetate (100 ml) and washed with 2M hydrochloric acid, brine, saturated aqueous sodium bicarbonate solution, brine and dried. The residue obtained on removal of the solvent was triturated with diethyl ether and filtered. The solid was dissolved in warm acetone, treated with activated charcoal and filtered. The solvent was removed to give Ac-Leu-Leu-(2-furyl)acetonitrile (0.64 g). Mp 235-242°C; m/z 391 (MH)⁺; NMR 0.88 (m, 12H), 1.51, (m, 6H), 1.83 (d, 3H), 4.29 (m, 2H), 6.23 (d, 1H), 6.53 (m, 2H), 7.76 (m, 1H), 7.95 (m, 2H), 9.19 (m, 1H).

Example 42**(2S)-2-(Boc-Leu-NH)isovaleronitrile**

Trifluoroacetic anhydride (0.84 g) was added dropwise to a mixture of BOC-Leu-Leu-NH₂ (1 g) and pyridine (10 ml) which was stirred under argon at -10°C. The mixture was

allowed to warm to room temperature over 1 hour, diluted with water and extracted with diethyl ether. The extract was washed successively with 1M hydrochloric acid and brine, dried and evaporated to dryness. The residue was recrystallized from ether/hexane to give (2S)-2-(BOC-Leu-NH)isovaleronitrile (0.66 g). Mp 117-118°C; m/z 326 (MH)⁺.

5

Example 43**(2S)-2-(Boc-Leu-NH)-4-methylthiobutyronitrile**

The process described in Example 42 was repeated using BOC-Leu-Met-NH₂ instead of BOC-Leu-Leu-NH₂ to give (2S)-2-(BOC-Leu-NH)-4-methylthiobutyronitrile. Mp 69-71°C; m/z 344 (MH)⁺.

10

Example 44**(2S)-2-[(N-4-chlorobenzyl-Leu)-Leu-NH]-4-methylthiobutyronitrile**

A mixture of (2S)-2-[H-Leu-Leu-NH]-4-methylthiobutyronitrile (90mg), ethanol (5ml) and 4-chlorobenzaldehyde was stirred at ambient temperature for 1 hour. Acetic acid (0.05ml) was added followed by sodium cyanoborohydride (50mg) and stirring was continued for a further 3 hours. Acetic acid (0.2ml) was added and the mixture was left at room temperature for 16 hours. The mixture was diluted with water and basified with sodium hydrogen carbonate then extracted with ethyl acetate. The extract was dried and evaporated to dryness and the residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent to give (2S)-2-[(N-4-chlorobenzyl-Leu)-Leu-NH]-4-methylthiobutyronitrile as a gum (102mg). The gum was dissolved in ethyl acetate and the solution was acidified with ethereal HCl. The mixture was evaporated to dryness and the residue was triturated with ethyl acetate and the insoluble solid collected to give (2S)-2-[(N-4-chlorobenzyl-Leu)-Leu-NH]-4-methylthiobutyronitrile hydrochloride (75mg). Mp 146-147°C; NMR 0.9 (m, 12H), 1.4-1.8 (m, 6H), 2.0-2.2 (m, 5H), 2.55(m, 2H), 3.6 (m, 1H), 3.7-4.2 (m, 2H), 4.35 (m, 1H), 4.85 (m, 1H), 7.5(m, 4H), 8.9-9.1 (m, 2H), 9.45 (m, 1H), 9.7 (m, 1H); m/z 481 (MH)⁺.

15

20


25

Example 45**2-[2-benzyloxycarbonylamino-2-phenylthioacetamido]-2-(2-thienyl)-acetonitrile**

A mixture of 2-benzyloxycarbonylamino-2-phenylthioacetic acid (160mg), 2-amino-2-(2-thienyl)acetonitrile hydrochloride (88 mg), hydroxybenzotriazole (75 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (105 mg) and 4-methylmorpholine (0.15 ml) in *N,N*-dimethylformamide (3 ml) was stirred at ambient temperature for 48 hours. The solvent was removed under reduced pressure and the residue was partitioned between ethyl acetate (100 ml) and water (100 ml). The ethyl acetate layer was separated and washed with 2M hydrochloric acid, brine, saturated aqueous sodium bicarbonate and dried and evaporated to dryness. The residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent followed by recrystallisation from a mixture of ethyl acetate and hexane to give 2-(2-benzyloxycarbonylamino-2-phenylthioacetamido)-2-(2-thienyl)-acetonitrile (4.53 g). M/z 438 (MH^+); NMR ($CDCl_3$) 5.1 (m, 2H), 5.5 (m, 1H), 5.85 (m, 1H), 6.2 (m, 1H), 6.8-7.1 (m, 2H), 7.2-7.55 (m, 12H).

Example 46 - 55

The following examples were prepared by a process similar to that described in Example 45:



Example	R^{11}	R^{10}	R^2	Mp ($^{\circ}C$)	MH^+
46	benzyloxy	2-propyl	H	-	322
47	benzyloxy	phenyl	H	-	356
48	benzyl	phenyl	H	-	340
49	2,4,6-trichloro phenoxyethyl	phenyl	H	153-154	458
50	naphth-2-yl	phenyl	H	151-153	376
51	2,4-dichlorophenyl	phenyl	H	171-172	394


52	naphth-2-yl	naphth-1-yl	H	190-192	426
53	naphth-2-yl	naphth-2-yl	H	186-188	426
54	phenylsulphonyl methyl	phenyl	H	189-190	404
55	benzyloxy	m-tolyl	thiophen-2- yl	118-121	452

Example 562-[2-benzyloxycarbonylamino-2-(4-chlorophenylthio)acetamido]-2-(2-thienyl)-acetonitrile

A mixture of 2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-
 5 acetonitrile (132 mg), 1,2-dichloroethane (5 ml), 2-naphthalenesulphonic acid (5 mg) and 4-
 chlorothiophenol (1.65 g) was stirred at reflux for 2 hours then cooled to room temperature
 and evaporated to dryness. The residue was dissolved in ethyl acetate and the solution was
 washed successively with 1M NaOH and brine, dried and evaporated to dryness. The residue
 was recrystallized from ethyl acetate and hexane to give 2-[2-benzyloxycarbonylamino-2-(4-
 10 chlorophenylthio)acetamido]-2-(2-thienyl)-acetonitrile (80 mg). Mp 165-166°C; m/z 390
 (MH)⁺; NMR 4.2 (d, 2H), 5.0 (q, 2 1H), 5.7 (d, 1H), 7.2-7.55 (m, 9H), 8.2 (d, 1H), 9.0 (t, 1H).

Examples 57-8

The following examples were prepared by a process similar to that described in
 15 Example 56:




Example	R ¹⁰	Mp (°C)	MH ⁺
57	3-chlorophenyl	149-150	390
58	benzyl	139-140	370

Example 59**N-(N-Morpholinocarbonyl)-(L)-leucyl-2-(2-thienyl)glycinenitrile**

A mixture of N-(N-morpholinocarbonyl)-(L)-leucine (Method M) (0.323 g), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (0.28 g) and 1-hydroxybenzotriazole (0.197 g) in N,N-dimethylformamide (50 ml) was stirred at ambient temperature for 30 minutes. 2-(2-Thienyl)-2-aminoacetonitrile (Method A1) (0.253 g) and N-methylmorpholine (160 μ l) were added and the mixture was stirred for 20 hours. The reaction mixture was evaporated to dryness (high vac) and the residue was suspended in saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate (2x50 ml). The combined ethyl acetate extracts were washed successively with 10% citric acid and brine and dried. The residue obtained on removal of the solvent was chromatographed on a Bond-elut column eluting with a mixture of ethyl acetate and dichloromethane (1/1 v/v) to give the title compound (0.159 g) as a 1/1 mixture of diastereoisomers. NMR: 9.2-9.3 (1H), 7.6 (m, 1H), 7.2 (m, 1H), 7.0 (m, 1H), 6.5 (m, 1H), 6.3 (m, 1H), 4.2 (m, 1H), 3.45 (m, 4H), 3.3 (m, 4H), 1.6 (m, 2H), 1.4 (m, 1H), 0.8 (m, 6H).

Examples 60-63

Following the method outlined in Example 59 and using the appropriate acid chlorides there were prepared:



20

Ex No.	R ¹¹	(M+H)
60	c-Hexyl	362
61	c-Pentyl	348
62	c-Hexylmethyl	376
63	N-Piperidiny	363

Example 64**N-[(2-Phenylacetyl-amino-2-phenylthio)acetyl]-2-thienylaminoacetonitrile**

Carbonyl diimidazole (0.538g) was added to a solution of N-[(2-phenylacetyl-amino-2-phenylthio)acetic acid (1g) in THF (25 ml) and the mixture was stirred at 20 °C for 20 hours.

- 5 2-(2-Thienyl)-aminoacetonitrile (Method A1) (0.579 g) and triethylamine (0.664 g) were added and the mixture was stirred at 20 °C for 20 hours. The solvent was removed and the residue dissolved in dichloromethane (25 ml) and washed successively with aqueous sodium bicarbonate solution (2x20 ml) and 2M hydrochloric acid (2x20 ml). The solvent was removed and the residue was chromatographed on silica eluting with a mixture of ethyl acetate and isohexane (35:100) to give the title compound.
- 10


Diastereoisomer 1; Faster running fraction: Mp 172 °C; m/z 422 (M+H)⁺; NMR: 9.9 (d, 1H), 8.95 (d, 1H), 7.65 (m, 1H), 7.2 (m, 11H), 6.45 (d, 1H), 5.8 (d, 1H), 3.5 (m, 2H).

Diastereoisomer 2; Slower running fraction: Mp 159 °C; m/z 422 (M+H)⁺; NMR: 9.8 (d, 1H), 8.95 (d, 1H), 7.65 (m, 1H), 7.2 (m, 11H), 6.45 (d, 1H), 5.8 (d, 1H), 3.5 (m, 2H).

15

Examples 65-66

Following the method outlined in Example 64 and using the appropriate starting materials there were prepared:



Ex No.	R ²	M.p. °C
65 ¹	2-thienyl	144
66	H	204

20 ¹ Mixture of diastereoisomers.

Example 67**N-[(2-Phenylacetyl-amino-2-phenylthio)acetyl]-2-thienylaminoacetonitrile**


Carbonyl diimidazole (0.538g) was added to a solution of N-[(2-phenylacetyl-amino-2-phenylthio)acetic acid (Method O) (1g) in THF (25 ml) and the mixture was stirred at 20 °C for 20 hours. 2-(2-Thienyl)-aminoacetonitrile (Method A1) (0.579 g) and triethylamine (0.664 g) were added and the mixture was stirred at 20 °C for 20 hours. The solvent was removed and the residue dissolved in dichloromethane (25 ml) and washed successively with aqueous sodium bicarbonate solution (2x20 ml) and 2M hydrochloric acid (2x20 ml). The solvent was removed and the residue was chromatographed on silica eluting with a mixture of ethyl acetate and isohexane (35:100) to give the title compound.

Diastereoisomer 1; Faster running fraction: Mp 172 °C; m/z 422 (M+H)⁺; NMR: 9.9 (d, 1H), 8.95 (d, 1H), 7.65 (m, 1H), 7.2 (m, 11H), 6.45 (d, 1H), 5.8 (d, 1H), 3.5 (m, 2H).

Diastereoisomer 2; Slower running fraction: Mp 159 °C; m/z 422 (M+H)⁺; NMR: 9.8 (d, 1H), 8.95 (d, 1H), 7.65 (m, 1H), 7.2 (m, 11H), 6.45 (d, 1H), 5.8 (d, 1H), 3.5 (m, 2H).

Examples 68-96

Following the method outlined in Example 67 and using the appropriate starting materials there were prepared:



Ex No.	R ²	R ¹⁰	R ¹¹	M.p. °C
68 ¹	2-thienyl	4-F-C ₆ H ₄	C ₆ H ₅ CH ₂	144
69	H	4-F-C ₆ H ₄	C ₆ H ₅ CH ₂	204
70 ²	2-thienyl	3-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	178
71 ³	2-thienyl	3-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	182
72	H	3-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	207
73	H	2-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	156
74	H	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	220
75	H	C ₆ H ₅ CH ₂	C ₆ H ₅ CH ₂	138

76	H	C ₆ H ₅ CH ₂ CH ₂	C ₆ H ₅ CH ₂	160
77	2-thienyl	C ₆ H ₅ CH ₂	C ₆ H ₅ CH ₂	174
78	2-thienyl	C ₆ H ₅ CH ₂ CH ₂	C ₆ H ₅ CH ₂	148
79	H	4-Cl-C ₆ H ₄	2-Cl-C ₆ H ₄ CH ₂	109
80	H	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄ CH ₂	178
81	H	4-Cl-C ₆ H ₄	4-Cl-C ₆ H ₄ CH ₂	190
82 ‘	2-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	181
83 ‘	2-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	166
84 “	4-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	192
85 “	4-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	168
86 ^	2-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄ CH ₂	185
87 ^	2-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄ CH ₂	183
88	3-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄ CH ₂	168
89	4-CH ₃ O-C ₆ H ₄	4-Cl-C ₆ H ₄	3-Cl-C ₆ H ₄ CH ₂	158
90 ~	H	4-Cl-C ₆ H ₄	<u>R</u> -C ₆ H ₅ CH(CH ₃)	151
91 ~	H	4-Cl-C ₆ H ₄	<u>R</u> -C ₆ H ₅ CH(CH ₃)	161
92 @	H	4-Cl-C ₆ H ₄	<u>S</u> -C ₆ H ₅ CH(CH ₃)	153
93 @	H	4-Cl-C ₆ H ₄	<u>S</u> -C ₆ H ₅ CH(CH ₃)	158
94 #	H	4-Cl-C ₆ H ₄	C ₆ H ₅ CH(CH ₂ CH ₃)	165
95 #	H	4-Cl-C ₆ H ₄	C ₆ H ₅ CH(CH ₂ CH ₃)	179
96	H	4-Cl-C ₆ H ₄	4-Cl-C ₆ H ₄ C(CH ₃) ₂	

¹ Mixture of diastereoisomers.

² Diastereoisomer 1; Faster running fraction.

³ Diastereoisomer 2; Slower running fraction.

*, ‘, “, ^, ~, @, # Pairs of diastereoisomers.

5 Example 97


2-[2-benzyloxycarbonylamino-2-phenylthioacetamido]-2-(2-thienyl)-acetonitrile

A mixture of 2-benzyloxycarbonylamino-2-phenylthioacetic acid (Method P) (160mg), 2-amino-2-(2-thienyl)acetonitrile hydrochloride (Method A1) (88 mg), hydroxybenzotriazole (75 mg), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (105 mg) and 4-methylmorpholine (0.15 ml) in *N,N*-dimethylformamide (3 ml) was stirred at ambient

temperature for 48 hours. The solvent was removed under reduced pressure and the residue was partitioned between ethyl acetate (100 ml) and water (100 ml). The ethyl acetate layer was separated and washed with 2M hydrochloric acid, brine, saturated aqueous sodium bicarbonate and dried and evaporated to dryness. The residue was purified by flash chromatography on silica (Merck, ART 9385) using increasingly polar mixtures of ethyl acetate and hexane as eluent followed by recrystallisation from a mixture of ethyl acetate and hexane to give 2-(2-benzyloxycarbonylamino-2-phenylthioacetamido)-2-(2-thienyl)-acetonitrile (4.53 g). M/z 438 (MH^+); NMR ($CDCl_3$) 5.1 (m, 2H), 5.5 (m, 1H), 5.85 (m, 1H), 6.2 (m, 1H), 6.8-7.1 (m, 2H), 7.2-7.55 (m, 12H).

Example 98-107

The following examples were prepared by a process similar to that described in Example 97:



Example	R ¹¹	R ¹⁰	R ²	Mp (°C)	MH ⁺
98	benzyloxy	2-propyl	H	-	322
99	benzyloxy	phenyl	H	-	356
100	benzyl	phenyl	H	-	340
101	2,4,6-trichloro phenoxyethyl	phenyl	H	153-154	458
102	naphth-2-yl	phenyl	H	151-153	376
103	2,4-dichlorophenyl	phenyl	H	171-172	394
104	naphth-2-yl	naphth-1-yl	H	190-192	426
105	naphth-2-yl	naphth-2-yl	H	186-188	426
106	phenylsulphonyl methyl	phenyl	H	189-190	404


107	benzyloxy	m-tolyl	thiophen-2-yl	118-121	452
-----	-----------	---------	---------------	---------	-----

Example 1082-[2-benzyloxycarbonylamino-2-(4-chlorophenylthio)acetamido]-2-(2-thienyl)-acetonitrile

- 5 A mixture of 2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-acetonitrile (Method R) (132 mg), 1,2-dichloroethane (5 ml), 2-naphthalenesulphonic acid (5 mg) and 4-chlorothiophenol (1.65 g) was stirred at reflux for 2 hours then cooled to room temperature and evaporated to dryness. The residue was dissolved in ethyl acetate and the solution was washed successively with 1M NaOH and brine, dried and evaporated to dryness.
- 10 The residue was recrystallized from ethyl acetate and hexane to give 2-[2-benzyloxycarbonylamino-2-(4-chlorophenylthio)acetamido]-2-(2-thienyl)-acetonitrile (80 mg). Mp 165-166°C; m/z 390 (MH)⁺; NMR 4.2 (d, 2H), 5.0 (q, 2 1H), 5.7 (d, 1H), 7.2-7.55 (m, 9H), 8.2 (d, 1H), 9.0 (t, 1H).

15 **Examples 109-110**

The following examples were prepared by a process similar to that described in Example 108:




Example	R ¹⁰	Mp (°C)	MH ⁺
109	3-chlorophenyl	149-150	390
110	benzyl	139-140	370

Example 111**N-[(2-Phenylacetyl-amino-2-{4-fluorophenylsulphonyl})acetyl] aminoacetonitrile**

m-Chloroperbenzoic acid (578 mg) was added to a suspension of N-[(2-Phenylacetyl-amino-2-{4-fluorophenylthio})acetyl]aminoacetonitrile (Example 69) (300 mg) in dichloromethane (30 ml) and the mixture was stirred at room temperature for 3 hours then washed successively with aqueous sodium bicarbonate (3 x 10 ml), aqueous sodium thiosulphate (1 x 10 ml) and the organic layer was collected and dried. The residue obtained on removal of the solvent was triturated with diethyl ether to give the title compound Mp 175°C; NMR 9.39 (t, 1H), 9.3 (d, 1H), 7.75 (m, 2H), 7.3 (m, 7H), 6.05 (d, 1H), 4.29 (d, 2H), 3.5 (q, 2H).

Examples 112-116

Following the method outlined in Example 111 and using the appropriate starting materials there were prepared:



Ex No.	R ²	R ¹⁰	R ¹¹	M.p. °C
112	H	3-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	202
113	H	4-Cl-C ₆ H ₄	C ₆ H ₅ CH ₂	210
114	H	C ₆ H ₅	C ₆ H ₅ CH ₂	192
115 **	H	4-Cl-C ₆ H ₄	<u>S</u> -C ₆ H ₅ CH(CH ₃)	185
116 **	H	4-Cl-C ₆ H ₄	<u>S</u> -C ₆ H ₅ CH(CH ₃)	188

** Pair of diastereomers

Preparation of Starting Materials

The starting materials for the Examples above are either commercially available or are readily prepared by standard methods from known materials. For example the following reactions are illustrations but not limitations of the preparation of some of the starting materials used in the above reactions.

Method A**2-(2-Furyl)-2-aminoacetonitrile**

Ammonium chloride (25 g) was added to a solution of 2-furfuraldehyde (25 g) in diethyl ether (250 ml). A solution of sodium cyanide (17 g) in water (80 ml) was added over 20 minutes. The reaction mixture was stirred at ambient temperature for 14 hours, the aqueous layer was removed and the organic layer was washed twice with saturated aqueous sodium hydrogen carbonate solution (100 ml each time), dried and evaporated to dryness. The residue was dissolved in diethyl ether (250 ml) and cooled to 0 °C. Hydrogen chloride gas was bubbled through the solution keeping the temperature below 10 °C. 2-(2-Furyl)-2-aminoacetonitrile hydrochloride was filtered and dried, yield 33 g. ¹H NMR 6.19 (s, 1H), 6.56 (m, 1H), 6.78 (d, 1H), 7.83 (m, 1H), 9.83 (broad s, 2H).

Method A1

Following the method outlined in Method A and using the appropriate aldehyde there was prepared:

A1 2-(2-thienyl)-2-aminoacetonitrile hydrochloride

Method B**Z-Leu-Leu-Phe-NH₂**

A mixture of H-Leu-Phe-NH₂ hydrochloride (0.8 g), Z-Leu-OH (0.72 g), *N,N*-dimethylformamide (10 ml), 1-hydroxybenzotriazole (0.41 g), 1-dimethylaminopropyl-3-ethylcarbodiimide hydrochloride (0.57 g) and *N*-methylmorpholine (1 ml) was stirred at room temperature for 18 hours. The mixture was evaporated to dryness and the residue stirred with water and ethyl acetate. The insoluble solid was collected to give Z-Leu-Leu-Phe-NH₂ (0.67 g). *M/z* 525 (MH)⁺.

Method B1

Following the method outlined in Method C and using the appropriate protected starting materials there was prepared:

Method	Product	Starting Materials	(MH) ⁺
B1	Z-Phe-Leu-Met-NH ₂	H-Leu-Met-NH ₂ and Z-Phe-OH	543

Method C**Boc-Leu-Met-NH₂**

A mixture of Leu-Met-NH₂ hydrochloride (0.45 g), BOC-Leu-OH (0.35 g), *N,N*-dimethylformamide (5 ml), 1-hydroxybenzotriazole (0.23 g), dicyclohexylcarbodiimide (0.35 g) and *N*-methylmorpholine (0.2 ml) was stirred at room temperature for 18 hours. The mixture was filtered and the filtrate was evaporated to dryness. The residue was dissolved in ethyl acetate. The solution was washed successively with 0.5 M hydrochloric acid, 1M sodium hydroxide and brine, dried and evaporated to dryness. The residue was recrystallized from a mixture of ethyl acetate and hexane to give BOC-Leu-Met-NH₂ (0.45 g). Mp 145-147°C; m/z 475 (MH)⁺.

Methods C1-2

Following the method outlined in Method C and using the appropriate protected amino acid there was prepared:

Method	Product	Amino Acid	(MH) ⁺
C1	Z-Leu-Leu-Met-NH ₂	Z-Leu-OH	509
C2	[(2RS)-N-benzoyl-3-(4-pyridyl)alaninyl]-Leu-Met-NH ₂	(2RS)-N-benzoyl-3-(4-pyridyl)alanine	509

Method D**2-(2-propylthio)-2-(Z-Leu-Phe-NH)acetic acid**

A mixture of Z-Leu-Phe-NH₂ (3 g), glyoxylic acid monohydrate (0.83 g) and dioxan (20 ml) was stirred at reflux for 18 hours, cooled and evaporated to dryness. The residue was treated with dichloroethane (20 ml), 2-propane thiol (2.25 g) and naphthalene-2-sulphonic acid (50 mg) and the mixture was stirred at 60°C for 4 hours and then evaporated to dryness. The residue was stirred with ether and the insoluble white solid was collected to give 2-(2-propylthio)-2-(Z-Leu-Phe-NH)acetic acid (2.1 g). Mp 156-158°C; m/z 544 (MH)⁺.

Methods D1-2

Following the method outlined in Method D and using the appropriate dipeptide there was prepared:

Method	Product	Dipeptide
D1	2-(2-propylthio)-2-(Z-Leu-Leu-NH)acetic acid	Z-Leu-Leu-NH ₂

D2	2-(2-propylthio)-2-(Z-Phe-Leu-NH)acetic acid	Z-Phe-Leu-NH ₂
----	--	---------------------------

Method E**2-(2-propylthio)-2-(Z-Leu-Phe-NH)acetamide**

A mixture of 2-(2-propylthio)-2-(Z-Leu-Phe-NH)acetic acid (2 g), ammonium chloride (0.54 g), *N,N*-dimethylformamide (15 ml), 1-hydroxybenzotriazole (0.67 g), 1-dimethylaminopropyl-3-ethylcarbodiimide (0.95 g) and *N*-methylmorpholine (2 ml) was stirred at room temperature for 18 hours. The mixture was evaporated to dryness and the residue taken up in ethyl acetate. The solution was washed successively with 0.5 M hydrochloric acid, 1 M sodium hydroxide and brine. The solid which separated from the ethyl acetate phase was collected to give 2-(2-propylthio)-2-(Z-Leu-Phe-NH)acetamide, (1 g).

Mp 198-201°C; m/z 543 (MH)⁺.

Methods E1-2

Following the method outlined in Method E and using the appropriate acetic acid there was prepared:

Method	Product	Acetic Acid	(MH) ⁺
E1	2-(2-propylthio)-2-(Z-Leu-Leu-NH)acetamide	2-(2-propylthio)-2-(Z-Leu-Leu-NH)acetic acid	509
E2	2-(2-propylthio)-2-(Z-Phe-Leu-NH)acetamide	2-(2-propylthio)-2-(Z-Phe-Leu-NH)acetic acid	543

Method F**Z-Leu-Cha-OH**

A mixture of Z-Leu-OH (2.77 g), (*S*)-3-cyclohexylalanine methyl ester hydrochloride (1.92 g), 1-hydroxybenzotriazole (1.76 g), 1-dimethylaminopropyl-3-ethylcarbodiimide (1.92 g), *N,N*-dimethylformamide (10 ml) and *N*-methylmorpholine (3 ml) was stirred at room temperature for 18 hours. The mixture was diluted with ethyl acetate, and the solution was washed successively with 0.5 M hydrochloric acid, 1M sodium hydroxide and brine, then dried and evaporated to dryness. The residue was stirred in a mixture of tetrahydrofuran (15 ml) and 1M sodium hydroxide for 1 hour at room temperature. The mixture was acidified with 2M hydrochloric acid and extracted with ethyl acetate. The ethyl acetate layer was washed with brine, dried and evaporated to dryness to give Z-Leu-Cha-OH. M/z 419 (MH)⁺.

Method G**(2S)-2-(H-Leu-Leu-NH)-4-methylthiobutyronitrile**

Iodotrimethylsilane (0.6 ml) was added dropwise to a stirred solution of (2S)-2-(BOC-Leu-Leu-NH)-4-methylthiobutyronitrile (0.65 g) in dichloromethane (10 ml) the mixture was stirred at room temperature for 1 hour and then evaporated to dryness. The residue was redissolved in dichloromethane and the solution was treated with methanol (1 ml) and stirred until effervescence ceased. The solution was washed with aqueous sodium bicarbonate solution, dried and evaporated to dryness to give (2S)-2-(H-Leu-Leu-NH)-4-methylthiobutyronitrile as a gum (0.45 g). M/z 357 (MH)⁺.

Method H**Boc-Leu-Leu-NH₂**

A mixture of BOC-Leu-OH (6.93 g), Leu-OMe hydrochloride (5.45 g), 1-hydroxybenzotriazole (4.73 g), 1-dimethylaminopropyl-3-ethylcarbodiimide (6.7 g), *N,N*-dimethylformamide (50 ml) and *N*-methylmorpholine (10 ml) was stirred at room temperature for 18 hours. The mixture was evaporated to dryness and the residue was partitioned between water and ethyl acetate. The organic phase was washed successively with 0.5M hydrochloric acid, 1M sodium hydroxide and brine and then dried and evaporated to dryness. The residue was triturated with a mixture of ether and hexane to give Boc-Leu-Leu-OMe (11 g).

A mixture of a portion of the above ester (4 g), methanol (30 ml) and concentrated aqueous ammonia (50 ml) was stirred at room temperature for 72 hours. The mixture was diluted with water and extracted with ethyl acetate. The extract was washed with brine, dried and evaporated to dryness to give Boc-Leu-Leu-NH₂ (3.3 g). M/z 344.5 (MH)⁺.

Method I**(2S)-2-[(N-phenylacetyl-N-methyl-Leu)-Leu-Met-NH₂]**

Phenylacetyl chloride (0.57 g) was added, dropwise to a stirred, ice cooled solution of *N*-methyl-Leu (0.53 g) in 2M sodium hydroxide (4 ml) and the mixture was stirred at ambient temperature for 3 hours. The mixture was washed with ether and the aqueous phase was acidified with 2M hydrochloric acid and the precipitate collected to give *N*-phenylacetyl-*N*-methyl-Leu-OH (0.33 g). Mp 146-147°C; m/z 264 (MH)⁺.

This was then coupled with H-Leu-Met-NH₂ according to method C to give (2S)-2-[(N-phenylacetyl-N-methyl-Leu)-Leu-Met-NH₂]. M/z 507 (MH)⁺.

Method I1

(2S)-2-[(N-benzoyl-N-methyl-Leu)-Leu-Met-NH₂ [m/z 493 (MH)⁺] was also obtained by the process described in Method H.


Method J5 **2-benzyloxycarbonylamino-2-phenylthioacetic acid**

A mixture of benzyl carbamate (15.1 g), ether (100ml) and glyoxylic acid monohydrate (10.1 g) was stirred at room temperature for 16 hours. The thick suspension was filtered and the residue was washed with a mixture of ether and hexane to give 2-benzyloxycarbonylamino glycollic acid (17 g), which was used without further purification.

10 A mixture of 2-benzyloxycarbonylamino glycollic acid (2.25 g), 1,2-dichloroethane (50ml) and thiophenol (1.65 g) was stirred at reflux for 2 hours then cooled to room temperature. The mixture was extracted twice with aqueous sodium hydrogen carbonate and the combined extracts were washed with ether then acidified with 2M hydrochloric acid. The mixture was extracted with ether and the extract was dried and evaporated to dryness to give
15 2-benzyloxycarbonylamino-2-phenylthioacetic acid as a white solid (2.45 g). M/z 318 (MH)⁺.

Method J 1 - 4

Following the method outlined in Method J and using the appropriate amide instead of benzyl carbamate in the first stage and the appropriate thiol in the second stage there was prepared:



20


Method	R ¹	R ²	MH ⁺
J1	benzyloxy	2-propyl	322
J2	2,4,6-trichlorophenoxyethyl	phenyl	420
J3	2,4-dichlorophenyl	phenyl	356
J4	benzyloxy	m-tolyl	332

Method K**2-phenylacetamido-2-phenylthioacetic acid**

A mixture of phenylacetamide (2.7 g), glyoxylic acid monohydrate (2.02 g), 1,2-dichloroethane (100 ml), thiophenol (3.3 g) and 2-naphthalenesulphonic acid (100 mg) was stirred at reflux for 16 hours. The mixture was cooled and extracted twice with aqueous sodium hydrogen carbonate and the combined extracts were washed with ether, acidified with 2M hydrochloric acid and extracted with ether. The ether extract was dried and evaporated to dryness to give 2-phenylacetamido-2-phenylthioacetic acid. M/z 302 (MH⁺).

Method K 1 - 4

Following the method outlined in Method K and using the appropriate starting materials there was prepared:



Method	R ¹	R ²	MH ⁺
K1	naphth-2-yl	phenyl	338
K2	naphth-2-yl	naphth-1-yl	388
K3	naphth-2-yl	naphth-2-yl	388
K4	phenylsulphonylmethyl	phenyl	366

Method L**2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-acetonitrile**

A mixture of 2-benzyloxycarbonylamino glycollic acid (1.125 g), 2-aminoacetonitrile hydrochloride (0.7 g), hydroxybenzotriazole (0.75 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.05 g) and 4-methylmorpholine (3 ml) in *N,N*-dimethylformamide (20 ml) was stirred at ambient temperature for 18 hours. The solvent was removed under reduced pressure and the residue was stirred with a mixture of ethyl acetate (100 ml) and 1M Hydrochloric acid and the insoluble solid collected to give 2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-acetonitrile. Mp 146-148°C; m/z 264 (MH)⁺.

Method M**N-(N-Morpholinocarbonyl)-(L)-leucine**

Lithium hydroxide solution (5.9 g in 40 ml water) was added to a solution of N-(N-morpholinocarbonyl)-(L)-leucine methyl ester (Method N) (7.3 g) in THF (40 ml) and the mixture was stirred for 20 hours. The residue obtained on removal of the solvent was suspended in water (150 ml) and washed with ethyl acetate (50 ml). The aqueous layer was acidified to pH 1 with 2M HCl and extracted with ethyl acetate (3x70 ml). The combined ethyl acetate extracts were washed with water (75 ml) and brine (75 ml) and passed through phase-separating paper. Removal of the solvent gave N-(N-morpholinocarbonyl)-(L)-leucine as an oil. NMR: 6.5 (d, 1H), 4.0 (m, 1H), 3.5 (m, 4H), 3.3 (m, 4H), 1.5 (m, 2H), 1.4 (m, 1H), 0.8 (m, 6H).

Method N**N-(N-morpholinocarbonyl)-(L)-leucine methyl ester**

Triethylamine (8.4 ml) was added dropwise to L-leucine methyl ester hydrochloride (5 g) in dichloromethane (50 ml) at 0 °C followed by a solution of 4-morpholine carbonyl chloride (5 g) in dichloromethane (10 ml) and the mixture was stirred at ambient temperature for 20 hours. The reaction mixture was diluted with dichloromethane (100 ml) and washed with water (100 ml). The organic layer was collected and washed with 2M HCl (50 ml), brine (50 ml) and passed through phase-separating paper. Removal of the solvent gave N-(N-morpholinocarbonyl)-(L)-leucine methyl ester as a solid, 7.3 g. NMR: 6.7 (d, 1H), 4.1 (m, 1H), 3.6 (s, 2H), 3.5 (m, 4H), 3.3 (m, 4H), 1.6 (m, 2H), 1.4 (m, 1H), 0.8 (m, 6H); m/z: 259 (M+H)⁺.

Method O**N-[(2-phenylacetyl-amino-2-phenylthio)acetic acid**

This compound was prepared from phenylacetamide, glyoxylic acid and thiophenol following the procedure described in Tetrahedron, 31, 863 1975.

Method O1

Following the method outlined in Method O and using the appropriate starting material there was prepared:

N-[(2-phenylacetyl-amino-2-(4-fluorophenylthio)acetic acid.


Method P**2-benzyloxycarbonylamino-2-phenylthioacetic acid**

A mixture of benzyl carbamate (15.1 g), ether (100ml) and glyoxylic acid monohydrate (10.1 g) was stirred at room temperature for 16 hours. The thick suspension was filtered and the residue was washed with a mixture of ether and hexane to give 2-benzyloxycarbonylamino glycollic acid (17 g), which was used without further purification.

A mixture of 2-benzyloxycarbonylamino glycollic acid (2.25 g), 1,2-dichloroethane (50ml) and thiophenol (1.65 g) was stirred at reflux for 2 hours then cooled to room temperature. The mixture was extracted twice with aqueous sodium hydrogen carbonate and the combined extracts were washed with ether then acidified with 2M hydrochloric acid. The mixture was extracted with ether and the extract was dried and evaporated to dryness to give 2-benzyloxycarbonylamino-2-phenylthioacetic acid as a white solid (2.45 g). M/z 318 (MH)⁺.

Method P 1 - 4

Following the method outlined in Method P and using the appropriate amide instead of benzyl carbamate in the first stage and the appropriate thiol in the second stage there was prepared:



Method	R ¹	R ²	MH ⁺
P1	benzyloxy	2-propyl	322
P2	2,4,6-trichlorophenoxymethyl	phenyl	420
P3	2,4-dichlorophenyl	phenyl	356
P4	benzyloxy	m-tolyl	332


Method Q**2-phenylacetamido-2-phenylthioacetic acid**

A mixture of phenylacetamide (2.7 g), glyoxylic acid monohydrate (2.02 g), 1,2-dichloroethane (100 ml), thiophenol (3.3 g) and 2-naphthalenesulphonic acid (100 mg) was stirred at reflux for 16 hours. The mixture was cooled and extracted twice with aqueous sodium hydrogen carbonate and the combined extracts were washed with ether, acidified with

2M hydrochloric acid and extracted with ether. The ether extract was dried and evaporated to dryness to give 2-phenylacetamido-2-phenylthioacetic acid. M/z 302 (MH^+).

Method Q 1 - 4

Following the method outlined in Method Q and using the appropriate starting materials there was prepared:



Method	R ¹	R ²	MH ⁺
Q1	naphth-2-yl	phenyl	338
Q2	naphth-2-yl	naphth-1-yl	388
Q3	naphth-2-yl	naphth-2-yl	388
Q4	phenylsulphonylmethyl	phenyl	366


Method R

2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-acetonitrile

A mixture of 2-benzyloxycarbonylamino glycollic acid (1.125 g), 2-aminoacetonitrile hydrochloride (0.7 g), hydroxybenzotriazole (0.75 g), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.05 g) and 4-methylmorpholine (3 ml) in *N,N*-dimethylformamide (20 ml) was stirred at ambient temperature for 18 hours. The solvent was removed under reduced pressure and the residue was stirred with a mixture of ethyl acetate (100 ml) and 1M Hydrochloric acid and the insoluble solid collected to give 2-[2-benzyloxycarbonylamino-2-(hydroxy)acetamido]-2-(2-thienyl)-acetonitrile. Mp 146-148°C; m/z 264 (MH^+).

CLAIMS

1. A compound of formula (I):



(I)

wherein

r is 0 or 1;

R^1 is hydrogen, optionally substituted benzyl where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy,

- 10 trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl, or R^1 is a group of formula
- 15 (II):



(II)

wherein R^5 is C_{1-6} alkyl (optionally substituted with an optionally substituted phenyl, an optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted

20 phenoxy, optionally substituted phenylsulphonyl, optionally substituted C_{3-12} cycloalkyl or Het), C_{1-6} alkoxy, optionally substituted phenyl, optionally substituted naphthyl, optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted C_{3-12} cycloalkyl, Het or optionally substituted phenyl C_{1-6} alkoxy; where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy,

25 trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkoxycarbonyl, mercapto,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl,

N-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R² is H, C₁₋₆alkyl [optionally substituted with one or more of hydroxy,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, R⁴, R⁴C₁₋₆alkylsulphanyl,

5 R⁴C₁₋₆alkylsulphinyl, R⁴C₁₋₆alkylsulphonyl], or R² is C₁₋₆alkoxy [optionally substituted with one or more of C₂₋₆alkenyl, C₂₋₆alkynyl, R⁴, R⁴C₂₋₆alkenyl, R⁴C₂₋₆alkynyl, Het and trifluoromethyl], or R² is C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxycarbonyl, carbamoyl,

N-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, R⁴, R⁴S, R⁴C₁₋₆alkylsulphanyl,

N-(R⁴C₁₋₆alkyl)carbamoyl, *N*-(HetC₁₋₆alkyl)carbamoyl, C₁₋₆alkanoylamino,

10 C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl or C₁₋₆alkylsulphonyl; R⁴ is an optionally substituted phenyl or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms, said optional substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino,

15 *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl,


N-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl,


N-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R³ is H or C₁₋₆alkyl;


20 (AA¹) and (AA²) are independently chosen from Ala, Arg, Cys, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr, Val,



Lys(CHO),




Arg(NO₂),




β-Ala,


66




Ser(Bzl),




Ph-Gly,




Nle,




Ser(O^tBu),




His(Bzl),




Met(O),



Cha,




His(Me),




Cit,


5




Tyr(^tBu),




Met(O₂),




Pyr-Ala



Phe(S),




Leu(S),




Phe(CH₂S),

10


67




Cy(S)-Gly,




Hetar(S)-Gly,



alk(S)-Gly or




Het(S)-Gly, and

Phe(CH₂CH₂)S;

- 5 wherein Ring A is C₃₋₁₂cycloalkyl; Ring B is a 5 or 6 membered heteroaryl ring; Ring C is Het; V is C₁₋₆alkyl excluding isopropyl; the nitrogen of the amino acid may optionally be alkylated with C₁₋₆alkyl; the phenyl group of Phe(S) and Rings A and B are optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, 10 *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl or *N,N*-(C₁₋₆alkyl)₂sulphamoyl; the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group; the sulphur moiety in 15 the α -position of the amino acid (AA³) may be optionally oxidised to form an -S(O)₂- or -S(O)- moiety; and Het is a fully saturated monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms; or a pharmaceutically acceptable salt thereof.

20

2. A compound as claimed in claim 1 having the formula (Ia):




wherein:


R⁷ is optionally substituted benzyl, optionally substituted phenoxyethyl, optionally substituted phenylsulphonylmethyl, optionally substituted benzyloxy, optionally substituted naphthyl, optionally substituted phenyl or t-butoxy where said optional substituents are chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R⁶ is hydrogen, optionally substituted phenyl or optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms; said optional substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;


(AA³) is selected from:




Phe(S),




Leu(S),




Phe(CH₂S),




Cy(S)-Gly,



Hetar(S)-Gly,



alk(S)-Gly and



Het(S)-Gly, and

Phe(CH₂CH₂)S;

wherein Ring A is C₃₋₁₂cycloalkyl, Ring B is a 5 or 6 membered heteroaryl ring, Ring C is Het, V is C₁₋₆alkyl excluding isopropyl; the nitrogen of the amino acid may

optionally be alkylated with C₁₋₆alkyl; the phenyl group of Phe(S) and Rings A and B may be optionally substituted with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino, *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl,

N-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl; the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group; the sulphur moiety in the α -position of the amino acid (AA³) may be optionally oxidised to form an -S(O)₂- or -S(O)- moiety; and


Het is a fully saturated monocyclic 5 - 8 membered heterocyclic ring, with up to 4 ring heteroatoms; or a pharmaceutically acceptable salt thereof.

3. A compound as claimed in claim 2 wherein R⁶ is hydrogen, optionally substituted phenyl or a 5 membered heteroaryl ring containing a maximum of four heteroatoms.

4. A compound as claimed in claim 2 or 3 wherein R⁷ is benzyl (optionally substituted with halo (such as chloro)), α -(C₁₋₄ alkyl)-benzyl (optionally substituted with halo (such as chloro)), α,α -di(C₁₋₄ alkyl)-benzyl (optionally substituted with halo (such as chloro)), optionally substituted phoxymethyl, phenylsulphonylmethyl, benzyloxy, naphthyl or optionally substituted phenyl where said optional substituents are chosen from one or more halo.


5. A compound as claimed in claim 2, 3 or 4 wherein (AA³) is Leu(S), Phe(S) optionally substituted with C₁₋₆alkyl or halo and wherein the phenyl group of Phe(S) may be fused to another phenyl group to form a naphthyl group or the sulphur moiety in the α-position of the amino acid (AA) may be optionally oxidised to form an -S(O)₂- or Phe(CH₂S).

6. A process for preparing a compound of formula (Ia) as claimed in claim 2 comprising:
a) coupling an acid of formula (IIIa):




(IIIa)

or a reactive derivative thereof, with an amine of formula (IVa):



(IVa)


- b) dehydrating a compound of formula (Va):



(Va)


under standard conditions; or,

- c) reacting an amine of formula (VIa):



(VIa)

with an acid of formula (VIIa):




71

(VIIa)

or a reactive derivative thereof;
 wherein R^6 , R^7 and AA^3 are as defined in claim 2.

5 7. A compound of formula (I):



(I)

wherein:

r is 0 or 1;

10 R^1 is optionally substituted benzyl where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxycarbonyl, mercapto,
 15 C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl)₂sulphamoyl or R^1 is a group of formula (II):



(II)

20 wherein R^5 is C_{1-6} alkyl (optionally substituted with an optionally substituted phenyl, an optionally substituted 5 or 6 membered heteroaryl ring, optionally substituted phenoxy or optionally substituted phenylsulphonyl), C_{1-6} alkoxy, optionally substituted phenyl, optionally substituted naphthyl, optionally substituted phenyl C_{1-6} alkoxy where said optional substituents are chosen from one or more of C_{1-6} alkyl, halo,
 25 trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, amino, C_{1-6} alkylamino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, nitro, carboxy, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, C_{1-6} alkoxycarbonyl, mercapto, C_{1-6} alkylsulphanyl, C_{1-6} alkylsulphinyl,

C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R² is H, C₁₋₆alkyl [optionally substituted with one or more of hydroxy,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, **R**⁴, **R**⁴C₁₋₆alkylsulphanyl,

5 **R**⁴C₁₋₆alkylsulphinyl, **R**⁴C₁₋₆alkylsulphonyl], or **R**² is C₁₋₆alkoxy [optionally substituted with one or more of C₂₋₆alkenyl, C₂₋₆alkynyl, **R**⁴, **R**⁴C₂₋₆alkenyl, **R**⁴C₂₋₆alkynyl, Het and trifluoromethyl], or **R**² is C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxycarbonyl, carbamoyl,

N-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, **R**⁴, **R**⁴S, **R**⁴C₁₋₆alkylsulphanyl,

N-(**R**⁴C₁₋₆alkyl)carbamoyl, *N*-(HetC₁₋₆alkyl)carbamoyl, C₁₋₆alkanoylamino,

10 C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl wherein **R**⁴ is an optionally substituted phenyl, or an optionally substituted 5 or 6 membered heteroaryl ring containing a maximum of four heteroatoms said optional substituents being chosen from one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino,

15 *N,N*-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto, C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl, *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl;

R³ is H or C₁₋₆alkyl; and

20 (**AA**¹) and (**AA**²) are independently chosen from Ala, Arg, Cys, Gly, His, Ile, Leu, Lys, Met, Phe, Ser, Thr, Trp, Tyr, Val, Lys(CHO), Arg(NO₂), β-Ala, Ser(Bzl), Ph-Gly, Nle, Ser(O^tBu), His(Bzl), Met(O), Cha, His(Me), Cit, Tyr(^tBu), Met(O₂), Pyr-Ala, Phe(S), Leu(S) or Phe(CH₂S); wherein the nitrogen of the amino acid may optionally be

alkylated with C₁₋₆alkyl and the phenyl group of Phe(S) may be optionally substituted
25 with one or more of C₁₋₆alkyl, halo, trifluoromethyl, hydroxy, trifluoromethoxy, cyano, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, amino, C₁₋₆alkylamino,

N,N-(C₁₋₆alkyl)₂amino, C₁₋₆alkanoylamino, nitro, carboxy, carbamoyl,

N-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkoxycarbonyl, mercapto,

C₁₋₆alkylsulphanyl, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, sulphamoyl,


30 *N*-(C₁₋₆alkyl)sulphamoyl and *N,N*-(C₁₋₆alkyl)₂sulphamoyl or the phenyl group may be fused to another phenyl group to form a naphthyl group; or a pharmaceutically acceptable salt thereof.

8. A compound of formula (I) as claimed in claim 7 wherein r is 0; and R^2 is furyl, pyrazolyl (optionally substituted with one or more of methyl and bromo), imidazolyl, 1,2,4-triazolyl, benzyl, 2-methylthioethyl, isopropylthio, methoxy, isopropoxy and 2-propynyloxy.

9. A compound of formula (I) as claimed in claim 1 wherein r is 0; and R^2 is thienyl.


10. A process for preparing a compound of formula (I) as claimed in claim 1 comprising:

a) coupling an acid of formula (III):




(III)

or a reactive derivative thereof, with an amine of formula (IV):



(IV)

b) dehydrating a compound of formula (V):



(V)


under standard conditions;

c) for compounds of formula (I) where $r = 1$, coupling an acid of formula (VI):



(VI)


or a reactive derivative thereof as defined hereinbefore, with an amine of formula (VII):



(VII)


74

d) for compounds of formula (I) where R^1 is a group of formula (II), reacting an amine of formula (VIII):



(VIII)


5 with an acid of formula (IX):



(IX)


or a reactive derivative thereof as defined hereinbefore; or

10 e) for compounds of formula (I) where R^1 is optionally substituted benzyl, reacting an amine of formula (X):



(X)

i) with a compound of formula (XI):




(XI)

15

where $(R)_n$ are optional substituents as defined above and L is a displaceable group; or

ii) with an aldehyde of formula (XII):




(XII)

20

where $(R)_n$ are optional substituents as defined above and L is a displaceable group;


wherein R^1 , R^2 , R^3 , AA^1 , AA^2 and r are as defined in claim 1.

11. A compound of formula (V):



5 wherein R^1 , R^2 , R^3 , AA^1 , AA^2 and r are as defined in claim 1.

12. A compound of formula (VIII):



10 wherein R^1 , R^2 , R^3 , AA^1 , AA^2 and r are as defined in claim 1.

13. A pharmaceutical composition comprising a compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, as claimed in claim 1, 2, 8 or 9 and a pharmaceutically acceptable diluent or carrier.

15

14. A compound of formula (I) or (Ia), or a pharmaceutically acceptable salt thereof, as claimed in claim 1, 2, 8 or 9 for use as a medicament

15. The use of a compound of formula (I) or (Ia) as claimed in claim 1, 2, 8 or 9, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the inhibition of a cysteine protease in a warm blooded animal.

20

16. The use of a compound of formula (I) or (Ia) as claimed in claim 1, 2, 8 or 9, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for use in the treatment of chronic obstructive pulmonary disease in a warm blooded animal.

25

17. A method of treating a Cathepsin L or Cathepsin S mediated disease state in mammals which comprises administering to a mammal in need of such treatment an effective

76

amount of a compound of formula (I) or (Ia) as claimed in claim 1, 2, 8 or 9, or a pharmaceutically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No PCT/GB 00/00529				
A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D307/68 C07C255/29 C07D333/24 C07K5/062 C07K5/078 C07K5/065 C07C323/60 C07C317/50 A61K31/275 A61K31/277 A61K31/341 A61K31/381 A61P11/00				
According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07C C07D C07K A61K A61P				
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched				
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) BEILSTEIN Data, WPI Data, PAJ, CHEM ABS Data				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category *	Citation of document, with indication, where appropriate, of the relevant passages			Relevant to claim No.
X	S. SUZUE, ET AL.: "Hepatic agents. I. Synthesis of aminoacyl (and hydroxyacyl) aminoacetoneitriles" CHEMICAL AND PHARMACEUTICAL BULLETIN, vol. 16, no. 8, August 1968 (1968-08), pages 1417-1432, XP002108053 Pharmaceutical Society of Japan, Tokyo, JP ISSN: 0009-2363 the whole document <div style="text-align: center; margin-top: 10px;">--- -/--</div>			1-4,6,7, 10-14
<div style="display: flex; justify-content: space-between;"> <input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>				
* Special categories of cited documents :				
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*Z* document member of the same patent family</p> </div> </div>				
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">14 June 2000</div>			Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">30/06/2000</div>	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016			Authorized officer <div style="text-align: center; font-weight: bold;">English, R</div>	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	A.R. KATRITZKY, ET AL.: "Benzotriazole-assisted synthesis of alpha-aminonitriles and a conceptually novel method for peptide elongation" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 6, June 1990 (1990-06), pages 1853-1857, XP002140042 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X compounds 5g-5m, 6g-6m ---	1,2,4,7, 11
X	GB 1 073 288 A (KYORIN SEIYAKU) 21 June 1967 (1967-06-21) the whole document ---	1,7,10, 12-14
X	S.A. THOMPSON, ET AL.: "Carboxy-modified amino acids and peptides as protease inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 29, no. 1, January 1986 (1986-01), pages 104-111, XP002140043 American Chemical Society, Washington, DC, US ISSN: 0022-2623 compounds 3b, 3d, 4b, 4d ---	1,7, 10-12
X	D.T. ELMORE, ET AL.: "Thioesters of amino acid derivatives as thioacylating agents in thiopeptide synthesis" JOURNAL OF THE CHEMICAL SOCIETY, PERKIN TRANSACTIONS 1, no. 5, May 1988 (1988-05), pages 1051-1055, XP002140044 Royal Society of Chemistry, Letchworth, GB ISSN: 0300-922X page 1051, right-hand column, line 10 - line 26 ---	1-3,6,7, 10
X	H. MOSER, ET AL.: "Poly(dipeptamidium)-Salze: Definition und Methoden zur präparativen Herstellung" HELVETICA CHIMICA ACTA, vol. 69, 1986, pages 1224-1262, XP002049723 Verlag Helvetica Chimica Acta, Basle, CH ISSN: 0018-019X compounds 3, 4a, 5 --- -/--	1-3,6,7, 10-12

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCT/GB 00/00529

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	D.J. BUTTLE, ET AL.: "Affinity purification of the novel cysteine proteinase papaya proteinase IV, and papain from papaya latex" BIOCHEMICAL JOURNAL, vol. 261, no. 2, 15 July 1989 (1989-07-15), pages 469-476, XP000913659 Portland Press, London, GB ISSN: 0264-6021 page 471, right-hand column ---	1
X	S. LIU, ET AL.: "Effects of ligand homologation and ligand reactivity on the apparent kinetic specificity of papain" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1250, 1995, pages 43-48, XP002140045 Elsevier Science, Amsterdam, NL ISSN: 0006-3002 compound 3a ---	1
X	C.A. LEWIS, ET AL.: "Thiohemiacetal formation by inhibitory aldehydes at the active site" BIOCHEMISTRY, vol. 16, no. 22, 1 November 1977 (1977-11-01), pages 4890-4895, XP002140046 American Chemical Society, Washington, DC, US ISSN: 0006-2960 table II, entries 8, 11 ---	1
P, X	WO 99 24460 A (NOVARTIS) 20 May 1999 (1999-05-20) the whole document -----	1-17

INTERNATIONAL SEARCH REPORT

International Application No. PCT/GB 00 00529

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-10 (partially), 11, 12 (partially)

The initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claims may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). For these reasons, a meaningful search over the whole breadth of the claims is impossible. Consequently, the search has been restricted to compounds for which a pharmaceutical use is indicated (particularly as cysteine protease inhibitors).

Claim 11 deals with synthetic intermediates which are amides of di- and tripeptides, a group of compounds which are well-known in the prior art so that it is impossible to determine which parts of the claim may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT). No attempt was made to search this claim, but where documents relevant to this claim were found during the search in relation to the other claims, this has been noted in the International Search Report.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/00529

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 1073288 A		BE 662788 A CH 456623 A FR 4312 M US 3467691 A US 3697577 A	17-08-1965 16-09-1969 10-10-1972
WO 9924460 A	20-05-1999	AU 1487399 A	31-05-1999